


CHAPTER 2


2.1

```
>> q0 = 12;R = 50;L = 5;C = 1e-4;  
>> t = linspace(0,.7);  
>> q = q0*exp(-R*t/(2*L)).*cos(sqrt(1/(L*C)-(R/(2*L))^2)*t);  
>> plot(t,q)
```



2.2

```
>> z = linspace(-4,4);  
>> f = 1/sqrt(2*pi)*exp(-z.^2/2);  
>> plot(z,f)  
>> xlabel('z')  
>> ylabel('frequency')
```



PROPRIETARY MATERIAL. © The McGraw-Hill Companies, Inc. All rights reserved. No part of this Manual may be displayed, reproduced or distributed in any form or by any means, without the prior written permission of the publisher, or used beyond the limited distribution to teachers and educators permitted by McGraw-Hill for their individual course preparation. If you are a student using this Manual, you are using it without permission.

2.3 (a)

```
>> t = linspace(5,29,5)
```

```
t =
     5     11     17     23     29
```

(b)

```
>> x = linspace(-3,4,8)
```

```
x =
    -3    -2    -1     0     1     2     3     4
```

2.4 (a)

```
>> v = -3:0.5:1
```

```
v =
   -3.0000  -2.5000  -2.0000  -1.5000  -1.0000  -0.5000   0   0.5000  1.0000
```

(b)

```
>> r = 8:-0.5:0
```

```
r =
Columns 1 through 6
     8.0000     7.5000     7.0000     6.5000     6.0000     5.5000
Columns 7 through 12
     5.0000     4.5000     4.0000     3.5000     3.0000     2.5000
Columns 13 through 17
     2.0000     1.5000     1.0000     0.5000         0
```

2.5

```
>> F = [11 12 15 9 12];
>> x = [0.013 0.020 0.009 0.010 0.012];
>> k = F./x
```

```
k =
   1.0e+003 *
     0.8462     0.6000     1.6667     0.9000     1.0000
```

```
>> U = .5*k.*x.^2
```


```
U =
     0.0715     0.1200     0.0675     0.0450     0.0720
```

```
>> max(U)
```

```
ans =
     0.1200
```

2.6

```
>> TF = 32:3.6:82.4;
>> TC = 5/9*(TF-32);
>> rho = 5.5289e-8*TC.^3-8.5016e-6*TC.^2+6.5622e-5*TC+0.99987;
>> plot(TC,rho)
```



2.7

```
>> A = [.035 .0001 10 2;
0.02 0.0002 8 1;
0.015 0.001 19 1.5;
0.03 0.0008 24 3;
0.022 0.0003 15 2.5]
```


```
A =
    0.0350    0.0001   10.0000    2.0000
    0.0200    0.0002    8.0000    1.0000
    0.0150    0.0010   19.0000    1.5000
    0.0300    0.0008   24.0000    3.0000
    0.0220    0.0003   15.0000    2.5000
```

```
>> U = sqrt(A(:,2))./A(:,1).*(A(:,3).*A(:,4)./(A(:,3)+2*A(:,4))).^(2/3)
```

```
U =
    0.3624
    0.6094
    2.5053
    1.6900
    1.1971
```


2.8

```
>> t = 10:10:60;
>> c = [3.4 2.6 1.6 1.3 1.0 0.5];
>> tf = 0:75;
>> cf = 4.84*exp(-0.034*tf);
>> plot(t,c,'s',tf,cf,':')
>> xlim([0 75])
```



2.9

```
>> t = 10:10:60;
>> c = [3.4 2.6 1.6 1.3 1.0 0.5];
>> tf = 0:70;
>> cf = 4.84*exp(-0.034*tf);
>> semilogy(t,c,'s',tf,cf,'--')
```



The result is a straight line. The reason for this outcome can be understood by taking the common logarithm of the function to give,


$$\log_{10} c = \log_{10} 4.84 - 0.034t \log_{10} e$$

Because $\log_{10} e = 0.4343$, this simplifies to the equation for a straight line,

$$\log_{10} c = 0.6848 - 0.0148t$$


2.10

```
>> v = 10:10:80;
>> F = [25 70 380 550 610 1220 830 1450];
>> vf = 0:90;
>> Ff = 0.2741*vf.^1.9842;
>> plot(v,F,'d',vf,Ff,':')
```




2.11

```
>> v = 10:10:80;
>> F = [25 70 380 550 610 1220 830 1450];
>> vf = 0:90;
>> Ff = 0.2741*vf.^1.9842;
>> loglog(v,F,'d',vf,Ff,':')
```



2.12

```
>> x = linspace(0,3*pi/2);
>> s = sin(x);
>> sf = x-x.^3/factorial(3)+x.^5/factorial(5)-x.^7/factorial(7);
>> plot(x,s,x,sf,'--')
```



2.13 (a)

```
>> m=[83.6 60.2 72.1 91.1 92.9 65.3 80.9];
>> vt=[53.4 48.5 50.9 55.7 54 47.7 51.1];
>> g=9.81; rho=1.225;
>> A=[0.454 0.401 0.453 0.485 0.532 0.474 0.486];
>> cd=g*m./vt.^2;
>> CD=2*cd/rho./A
```

```
CD =
    1.0343    1.0222    0.9839    0.9697    0.9591    0.9698    1.0210
```

(b)

```
>> CDavg=mean(CD),CDmin=min(CD),CDmax=max(CD)
```


```
CDavg =
    0.9943
```

```
CDmin =
    0.9591
```

```
CDmax =
    1.0343
```


(b)

```
>> subplot(1,2,1);plot(m,A,'o')
>> xlabel('mass (kg)');ylabel('area (m^2)')
>> title('area versus mass')
>> subplot(1,2,2);plot(m,CD,'o')
>> xlabel('mass (kg)');ylabel('CD')
>> title('dimensionless drag versus mass')
```



2.14 (a)

```
t = 0:pi/50:10*pi;
subplot(2,1,1);plot(exp(-0.1*t).*sin(t),exp(-0.1*t).*cos(t))
title('(a)')
subplot(2,1,2);plot3(exp(-0.1*t).*sin(t),exp(-0.1*t).*cos(t),t);
title('(b)')
```

**2.15 (a)**

```
>> x = 2;
>> x ^ 3;
>> y = 8 - x
```

```
y =
     6
```


(b)

```
>> q = 4:2:10;
>> r = [7 8 4; 3 6 -2];
>> sum(q) * r(2, 3)
```

```
ans =
    -56
```

2.16

```
>> y0=0;v0=30;g=9.81;
>> x=0:5:100;
>> theta0=15*pi/180;
>> y1=tan(theta0)*x-g/(2*v0^2*cos(theta0)^2)*x.^2+y0;
>> theta0=30*pi/180;
>> y2=tan(theta0)*x-g/(2*v0^2*cos(theta0)^2)*x.^2+y0;
>> theta0=45*pi/180;
>> y3=tan(theta0)*x-g/(2*v0^2*cos(theta0)^2)*x.^2+y0;
>> theta0=60*pi/180;
>> y4=tan(theta0)*x-g/(2*v0^2*cos(theta0)^2)*x.^2+y0;
>> theta0=75*pi/180;
>> y5=tan(theta0)*x-g/(2*v0^2*cos(theta0)^2)*x.^2+y0;
>> y=[y1' y2' y3' y4' y5']
>> plot(x,y)
>> axis([0 100 0 50])
>> legend('15', '30', '45', '60', '75')
```


2.17

```
>> R=8.314;E=1e5;A=7E16;
>> Ta=273:5:333;
>> k=A*exp(-E./(R*Ta))
```

k =

```
Columns 1 through 10
    0.0051    0.0113    0.0244    0.0510    0.1040    0.2070    0.4030
    0.7677    1.4326    2.6213
Columns 11 through 13
    4.7076    8.3048   14.4030
```

```
>> subplot(1,2,1);plot(Ta,k)
>> subplot(1,2,2);semilogy(1./Ta,k)
```



The result in (b) is a straight line. The reason for this outcome can be understood by taking the common logarithm of the function to give,

$$\log_{10} k = \log_{10} A - \left(\frac{E}{R} \log_{10} e \right) \frac{1}{T_a}$$

Thus, a plot of $\log_{10} k$ versus $1/T_a$ is linear with a slope of $-(E/R)\log_{10} e = -5.2237 \times 10^3$ and an intercept of $\log_{10} A = 16.8451$.

PROPRIETARY MATERIAL. © The McGraw-Hill Companies, Inc. All rights reserved. No part of this Manual may be displayed, reproduced or distributed in any form or by any means, without the prior written permission of the publisher, or used beyond the limited distribution to teachers and educators permitted by McGraw-Hill for their individual course preparation. If you are a student using this Manual, you are using it without permission.