

## EXERCISES 2.1

## Exercises 2.1 (page 201)

- |   |   |   |                        |
|---|---|---|------------------------|
| 1. function   | 2. relation   | 3. domain   | 4. range               |
| 5. $y = f(x)$   | 6. $x$  | 7. dependent  | 8. difference quotient |
| 9. $D = \{2, 3, 4, 5\}; R = \{3, 4, 5, 6\}$<br>Each element of the domain is paired with only one element of the range. Function.   |   | 10. $D = \{5, 6, 7, 8\}; R = \{4\}$<br>Each element of the domain is paired with only one element of the range. Function.             |                        |
| 11. $D = \{1, 2, -5\}; R = \{3, 4, 5, 2\}$<br>1 is both paired with 3 and 4. Not a function.  |   | 12. $D = \{-1, 2, 0\}; R = \{2, -1, 1, 3\}$<br>0 is both paired with 1 and 3. Not a function.   |                        |
| 13. $\{(LSU, Tigers), (Georgia, Bulldogs), (MSU, Bulldogs), (Auburn, Tigers)\}$<br>$D = \{LSU, Georgia, MSU, Auburn\}; R = \{\text{Tigers, Bulldogs}\}$<br>Each element of the domain is paired with only one element of the range. Function.   |   |   |                        |
| 14. $\{(Jackson, Louisiana), (Jackson, Mississippi), (Jackson, Tennessee), (Alexandria, Virginia)\}$<br>$D = \{Jackson, Alexandria\}; R = \{\text{Louisiana, Mississippi, Tennessee, Virginia}\}$<br>Jackson is paired with Louisiana, Mississippi, and Tennessee. Not a function.  |   |   |                        |
| 15. $\{(76, \text{September 9}), (76, \text{October 12}), (78, \text{May 10}), (80, \text{June 1})\}$<br>$D = \{76, 78, 80\}; R = \{\text{September 9, October 12, May 10, June 1}\}$<br>76 is paired with September 9 and October 12. Not a function.  |   |   |                        |
| 16. $\{(\text{Architect}, \$73,090), (\text{Dentist}, \$149,310), (\text{Microbiologist}, \$66,260), (\text{Actuary}, \$93,680)\}$<br>$D = \{\text{Architect, Dentist, Microbiologist, Actuary}\}; R = \{\$73,090, \$149,310, \$66,260, \$93,680\}$<br>Each element of the domain is paired with only one element of the range. Function. |   |   |                        |
| 17. $y = x$<br>Each value of $x$ is paired with only one value of $y$ .<br><b>function</b>  | 18. $y - 2x = 0$<br>$y = 2x$<br>Each value of $x$ is paired with only one value of $y$ .<br><b>function</b> | 19. $y^2 = x$<br>$y = \pm \sqrt{x}$<br>At least one value of $x$ is paired with more than one value of $y$ .<br><b>not a function</b> |                        |
| 20. $y^2 - 4x = 1$<br>$y^2 = 4x + 1$<br>$y = \pm \sqrt{4x + 1}$<br>At least one value of $x$ is paired with more than one value of $y$ .<br><b>not a function</b>   | 21. $y = x^2$<br>Each value of $x$ is paired with only one value of $y$ .<br><b>function</b>                | 22. $y + 1 = 5x^3$<br>Each value of $x$ is paired with only one value of $y$ .<br><b>function</b>                                     |                        |

## EXERCISES 2.1

23.  $|y| = x$

$$y = \pm x$$

At least one value of  $x$  is paired with more than one value of  $y$ . **not a function**

26.  $y - |x| = 3$

$$y = |x| + 3$$

Each value of  $x$  is paired with only one value of  $y$ . **function**

24.  $2|y| = x - 4$

$$|y| = \frac{x-4}{2}$$

$$y = \pm \frac{x-4}{2}$$

At least one value of  $x$  is paired with more than one value of  $y$ . **not a function**

25.  $|x - 2| = y$

$$y = |x - 2|$$

Each value of  $x$  is paired with only one value of  $y$ . **function**

27.  $|x| = |y|$

$$|y| = |x|$$

$$y = \pm |x|$$

At least one value of  $x$  is paired with more than one value of  $y$ . **not a function**

28.  $|y| = |x - 2|$

$$y = \pm (x - 2)$$

At least one value of  $x$  is paired with more than one value of  $y$ . **not a function**

29.  $y = 7$ ; Each value of  $x$  is paired with only one value of  $y$ . **function**

30.  $x = 7$ ; At least one value of  $x$  is paired with more than one value of  $y$ . **not a function**

31.  $y - 7 = \sqrt{x}$

$$y = \sqrt{x} + 7$$

Each value of  $x$  is paired with only one value of  $y$ . **function**

32.  $y - \sqrt[3]{x} = 8$

$$y = \sqrt[3]{x} + 8$$

Each value of  $x$  is paired with only one value of  $y$ . **function**

33.  $x^3 + y^2 = 25$

$$y^2 = -x^3 + 25$$

$$y = \pm \sqrt{-x^3 + 25}$$

At least one value of  $x$  is paired with more than one value of  $y$ . **not a function**

34.  $(x - 1)^2 + y^2 = 16$

$$y^2 = 16 - (x - 1)^2$$

$$y = \pm \sqrt{16 - (x - 1)^2}$$

At least one value of  $x$  is paired with more than one value of  $y$ . **not a function**

35.  $f(x) = 3x + 5 \Rightarrow \text{domain} = (-\infty, \infty)$

36.  $f(x) = -5x + 2 \Rightarrow \text{domain} = (-\infty, \infty)$

37.  $f(x) = x^2 - x + 1 \Rightarrow \text{domain} = (-\infty, \infty)$

38.  $f(x) = x^3 - 3x + 2 \Rightarrow \text{domain} = (-\infty, \infty)$

39.  $f(x) = \sqrt{x - 2} \Rightarrow x - 2 \geq 0$

$$\text{domain} = [2, \infty)$$

40.  $f(x) = \sqrt{2x + 3} \Rightarrow 2x + 3 \geq 0$

$$\text{domain} = \left[-\frac{3}{2}, \infty\right)$$

41.  $f(x) = \sqrt{4 - x} \Rightarrow 4 - x \geq 0$

$$\text{domain} = (-\infty, 4]$$

42.  $f(x) = 3\sqrt{2 - x} \Rightarrow 2 - x \geq 0$

$$\text{domain} = (-\infty, 2]$$

43.  $f(x) = \sqrt{x^2 - 1} \Rightarrow x^2 - 1 \geq 0$

$$\text{domain} = (-\infty, -1] \cup [1, \infty)$$

44.  $f(x) = \sqrt{x^2 - 2x - 3} \Rightarrow x^2 - 2x - 3 \geq 0$

$$\text{domain} = (-\infty, -1] \cup [3, \infty)$$

**EXERCISES 2.1**

**45.**  $f(x) = \sqrt[3]{x+1} \Rightarrow \text{domain} = (-\infty, \infty)$       **46.**  $f(x) = \sqrt[3]{5-x} \Rightarrow \text{domain} = (-\infty, \infty)$

**47.**  $f(x) = \frac{3}{x+1} \Rightarrow x \neq -1$   
 $\text{domain} = (-\infty, -1) \cup (-1, \infty)$

**48.**  $f(x) = \frac{-7}{x+3} \Rightarrow x \neq -3$   
 $\text{domain} = (-\infty, -3) \cup (-3, \infty)$

**49.**  $f(x) = \frac{x}{x-3} \Rightarrow x \neq 3$   
 $\text{domain} = (-\infty, 3) \cup (3, \infty)$

**50.**  $f(x) = \frac{x+2}{x-1} \Rightarrow x \neq 1$   
 $\text{domain} = (-\infty, 1) \cup (1, \infty)$

**51.**  $f(x) = \frac{x}{x^2-4} = \frac{x}{(x+2)(x-2)}$   
 $x \neq -2, x \neq 2$   
 $\text{domain} = (-\infty, -2) \cup (-2, 2) \cup (2, \infty)$

**52.**  $f(x) = \frac{2x}{x^2-9} = \frac{2x}{(x+3)(x-3)}$   
 $x \neq -3, x \neq 3$   
 $\text{domain} = (-\infty, -3) \cup (-3, 3) \cup (3, \infty)$

**53.**  $f(x) = \frac{1}{x^2-4x-5} = \frac{1}{(x+1)(x-5)}$   
 $x \neq -1, x \neq 5$   
 $\text{domain} = (-\infty, -1) \cup (-1, 5) \cup (5, \infty)$

**54.**  $f(x) = \frac{x}{2x^2-16x+30} = \frac{x}{2(x-3)(x-5)}$   
 $x \neq 3, x \neq 5$   
 $\text{domain} = (-\infty, 3) \cup (3, 5) \cup (5, \infty)$

**55.**  $f(x) = |x| + 3 \Rightarrow \text{domain} = (-\infty, \infty)$       **56.**  $f(x) = 2|x-1| \Rightarrow \text{domain} = (-\infty, \infty)$

**57.**  $f(x) = 3x - 2$

$\begin{aligned} f(2) &= 3(2) - 2 \\ &= 6 - 2 \\ &= 4 \end{aligned}$	$\begin{aligned} f(-3) &= 3(-3) - 2 \\ &= -9 - 2 \\ &= -11 \end{aligned}$	$\begin{aligned} f(k) &= 3k - 2 \\ &= 3k^2 - 3 - 2 \\ &= 3k^2 - 5 \end{aligned}$
--	---	--

**58.**  $f(x) = 5x + 7$

$\begin{aligned} f(2) &= 5(2) + 7 \\ &= 10 + 7 \\ &= 17 \end{aligned}$	$\begin{aligned} f(-3) &= 5(-3) + 7 \\ &= -15 + 7 \\ &= -8 \end{aligned}$	$\begin{aligned} f(k) &= 5k + 7 \\ &= 5(k^2 - 1) + 7 \\ &= 5k^2 - 5 + 7 \\ &= 5k^2 + 2 \end{aligned}$
--	---	---

**59.**  $f(x) = \frac{1}{2}x + 3$

$\begin{aligned} f(2) &= \frac{1}{2}(2) + 3 \\ &= 1 + 3 \\ &= 4 \end{aligned}$	$\begin{aligned} f(-3) &= \frac{1}{2}(-3) + 3 \\ &= -\frac{3}{2} + 3 \\ &= \frac{3}{2} \end{aligned}$	$\begin{aligned} f(k) &= \frac{1}{2}k + 3 \\ &= \frac{1}{2}k^2 - \frac{1}{2} + 3 \\ &= \frac{1}{2}k^2 + \frac{5}{2} \end{aligned}$
--	---	--

**60.**  $f(x) = \frac{2}{3}x + 5$

$\begin{aligned} f(2) &= \frac{2}{3}(2) + 5 \\ &= \frac{4}{3} + 5 \\ &= \frac{19}{3} \end{aligned}$	$\begin{aligned} f(-3) &= \frac{2}{3}(-3) + 5 \\ &= -2 + 5 \\ &= 3 \end{aligned}$	$\begin{aligned} f(k) &= \frac{2}{3}k + 5 \\ &= \frac{2}{3}k^2 - \frac{2}{3} + 5 \\ &= \frac{2}{3}k^2 + \frac{13}{3} \end{aligned}$
---	---	---

**EXERCISES 2.1**

**61.**

$$f(x) = x^2$$

$f(2) = 2^2$	$f(-3) = (-3)^2$	$f(k) = k^2$	$f(k^2 - 1) = (k^2 - 1)^2$
$= 4$	$= 9$		$= (k^2 - 1)(k^2 - 1)$
			$= k^4 - 2k^2 + 1$

**62.**

$$f(x) = 3 - x^2$$

$f(2) = 3 - 2^2$	$f(-3) = 3 - (-3)^2$	$f(k) = 3 - k^2$	$f(k^2 - 1) = 3 - (k^2 - 1)^2$
$= 3 - 4$	$= 3 - 9$		$= 3 - (k^2 - 1)(k^2 - 1)$
$= -1$	$= -6$		$= 3 - (k^4 - 2k^2 + 1)$

$f(k^2 - 1) = 3 - (k^2 - 1)^2$	$= 3 - (k^4 - 2k^2 + 1)$	$= -k^4 + 2k^2 + 2$
--------------------------------	--------------------------	---------------------

**63.**

$$f(x) = x^2 + 3x - 1$$

$f(2) = 2^2 + 3(2) - 1$	$f(-3) = (-3)^2 + 3(-3) - 1$	$f(k) = k^2 + 3k - 1$
$= 4 + 6 - 1$	$= 9 - 9 - 1$	
$= 9$	$= -1$	

$$\begin{aligned} f(k^2 - 1) &= (k^2 - 1)^2 + 3(k^2 - 1) - 1 \\ &= k^4 - 2k^2 + 1 + 3k^2 - 3 - 1 \\ &= k^4 + k^2 - 3 \end{aligned}$$

**64.**

$$f(x) = -x^2 - 2x + 1$$

$f(2) = -(2)^2 - 2(2) + 1$	$f(-3) = -(-3)^2 - 2(-3) + 1$	$f(k) = -k^2 - 2k + 1$
$= -4 - 4 + 1$	$= -9 + 6 + 1$	
$= -7$	$= -2$	

$$\begin{aligned} f(k^2 - 1) &= -(k^2 - 1)^2 - 2(k^2 - 1) + 1 \\ &= -(k^4 - 2k^2 + 1) - 2k^2 + 2 + 1 \\ &= -k^4 + 2k^2 - 1 - 2k^2 + 3 \\ &= -k^4 + 2 \end{aligned}$$

**65.**

$$f(x) = x^3 - 2$$

$f(2) = 2^3 - 2$	$f(-3) = (-3)^3 - 2$	$f(k) = k^3 - 2$	$f(k^2 - 1) = (k^2 - 1)^3 - 2$
$= 8 - 2$	$= -27 - 2$		$= (k^2 - 1)(k^2 - 1)(k^2 - 1) - 2$
$= 6$	$= -29$		$= k^6 - 3k^4 + 3k^2 - 1 - 2$

$f(k^2 - 1) = (k^2 - 1)^3 - 2$	$= (k^2 - 1)(k^2 - 1)(k^2 - 1) - 2$	$= k^6 - 3k^4 + 3k^2 - 1 - 2$	$= k^6 - 3k^4 + 3k^2 - 3$
--------------------------------	-------------------------------------	-------------------------------	---------------------------

**EXERCISES 2.1**

**66.**

$$f(x) = -x^3$$

$f(2) = -(2^3)$ = -8	$f(-3) = -(-3)^3$ = -(-27) = 27	$f(k) = -(k)^3$ = $-k^3$	$f(k^2 - 1) = -(k^2 - 1)^3$ = $-(k^2 - 1)(k^2 - 1)(k^2 - 1)$ = $-(k^6 - 3k^4 + 3k^2 - 1)$ = $-k^6 + 3k^4 - 3k^2 + 1$
-------------------------	---------------------------------------	-----------------------------	---

**67.**

$$f(x) = |x^2 + 1|$$

$f(2) =  2^2 + 1 $ =  5  = 5	$f(-3) =  (-3)^2 + 1 $ =  10  = 10	$f(k) =  k^2 + 1 $ = $k^2 + 1$ $[k^2 + 1 \geq 0]$	$f(k^2 - 1) =  (k^2 - 1)^2 + 1 $ = $(k^2 - 1)^2 + 1$ = $k^4 - 2k^2 + 1 + 1$ = $k^4 - 2k^2 + 2$ $[(k^2 - 1)^2 + 1 \geq 0]$
------------------------------------	--	---	---

**68.**

$$f(x) = |x^2 + x + 4|$$

$f(2) =  2^2 + 2 + 4 $ =  4 + 2 + 4  =  10  = 10	$f(-3) =  (-3)^2 + (-3) + 4 $ =  9 - 3 + 4  =  10  = 10	$f(k) =  k^2 + k + 4 $ = $k^2 + k + 4$ $[k^2 + k + 4 \geq 0]$
---	--	---

$$\begin{aligned} f(k^2 - 1) &= |(k^2 - 1)^2 + (k^2 - 1) + 4| \\ &= |(k^2 - 1)^2 + k^2 + 3| \\ &= (k^2 - 1)^2 + k^2 + 3 \\ &= k^4 - 2k^2 + 1 + k^2 + 3 \\ &= k^4 - k^2 + 4 \\ &\quad [(k^2 - 1)^2 + k^2 + 3 \geq 0] \end{aligned}$$

**69.**

$$f(x) = \frac{2}{x+4}$$

$f(2) = \frac{2}{2+4}$ = $\frac{2}{6} = \frac{1}{3}$	$f(-3) = \frac{2}{-3+4}$ = $\frac{2}{1} = 2$	$f(k) = \frac{2}{k+4}$	$f(k^2 - 1) = \frac{2}{k^2 - 1 + 4}$ = $\frac{2}{k^2 + 3}$
---	---	------------------------	---

**EXERCISES 2.1**

**70.**  $f(x) = \frac{3}{x-5}$

$f(2) = \frac{3}{2-5} = \frac{3}{-3} = -1$	$f(-3) = \frac{3}{-3-5} = \frac{3}{-8} = -\frac{3}{8}$	$f(k) = \frac{3}{k-5}$	$f(k^2 - 1) = \frac{3}{k^2 - 1 - 5} = \frac{3}{k^2 - 6}$
--	--	------------------------	--

**71.**  $f(x) = \frac{1}{x^2 - 1}$

$f(2) = \frac{1}{2^2 - 1} = \frac{1}{4 - 1} = \frac{1}{3}$	$f(-3) = \frac{1}{(-3)^2 - 1} = \frac{1}{9 - 1} = \frac{1}{8}$	$f(k) = \frac{1}{k^2 - 1}$	$f(k^2 - 1) = \frac{1}{(k^2 - 1)^2 - 1} = \frac{1}{k^4 - 2k^2 + 1 - 1} = \frac{1}{k^4 - 2k^2}$
--	--	----------------------------	--

**72.**  $f(x) = \frac{3}{x^2 + 3}$

$f(2) = \frac{3}{2^2 + 3} = \frac{3}{4 + 3} = \frac{3}{7}$	$f(-3) = \frac{3}{(-3)^2 + 3} = \frac{3}{9 + 3} = \frac{3}{12} = \frac{1}{4}$	$f(k) = \frac{3}{k^2 + 3}$	$f(k^2 - 1) = \frac{3}{(k^2 - 1)^2 + 3} = \frac{3}{k^4 - 2k^2 + 1 + 3} = \frac{3}{k^4 - 2k^2 + 4}$
--	---	----------------------------	--

**73.**  $f(x) = \sqrt{x^2 + 1}$

$f(2) = \sqrt{2^2 + 1} = \sqrt{4 + 1} = \sqrt{5}$	$f(-3) = \sqrt{(-3)^2 + 1} = \sqrt{9 + 1} = \sqrt{10}$	$f(k) = \sqrt{k^2 + 1}$	$f(k^2 - 1) = \sqrt{(k^2 - 1)^2 + 1} = \sqrt{k^4 - 2k^2 + 1 + 1} = \sqrt{k^4 - 2k^2 + 2}$
---	--	-------------------------	---

**74.**  $f(x) = \sqrt{x^2 - 1}$

$f(2) = \sqrt{2^2 - 1} = \sqrt{4 - 1} = \sqrt{3}$	$f(-3) = \sqrt{(-3)^2 - 1} = \sqrt{9 - 1} = \sqrt{8} = 2\sqrt{2}$	$f(k) = \sqrt{k^2 - 1}$	$f(k^2 - 1) = \sqrt{(k^2 - 1)^2 - 1} = \sqrt{k^4 - 2k^2 + 1 - 1} = \sqrt{k^4 - 2k^2} = \sqrt{k^2(k^2 - 2)} =  k \sqrt{k^2 - 2}$
---	---	-------------------------	---

**75.**  $f(x) = \sqrt[3]{x} - 1$

$f(2) = \sqrt[3]{2} - 1$	$f(-3) = \sqrt[3]{-3} - 1 = -\sqrt[3]{3} - 1$	$f(k) = \sqrt[3]{k} - 1$	$f(k^2 - 1) = \sqrt[3]{k^2 - 1} - 1$
--------------------------	---	--------------------------	--------------------------------------

**EXERCISES 2.1**

- 76.**  $f(x) = \sqrt[3]{x+1}$
- |                        |                          |                        |                                      |
|------------------------|--------------------------|------------------------|--------------------------------------|
| $f(2) = \sqrt[3]{2+1}$ | $f(-3) = \sqrt[3]{-3+1}$ | $f(k) = \sqrt[3]{k+1}$ | $f(k^2 - 1) = \sqrt[3]{k^2 - 1 + 1}$ |
| $= \sqrt[3]{3}$        | $= \sqrt[3]{-2}$         |                        | $= \sqrt[3]{k^2}$                    |
| $= -\sqrt[3]{2}$       |                          |                        |                                      |
- 77.** 
$$\begin{aligned} \frac{f(x+h) - f(x)}{h} &= \frac{[3(x+h)+1] - [3x+1]}{h} = \frac{[3x+3h+1] - [3x+1]}{h} \\ &= \frac{3x+3h+1 - 3x-1}{h} = \frac{3h}{h} = 3 \end{aligned}$$
- 78.** 
$$\begin{aligned} \frac{f(x+h) - f(x)}{h} &= \frac{[5(x+h)-1] - [5x-1]}{h} = \frac{[5x+5h-1] - [5x-1]}{h} \\ &= \frac{5x+5h-1 - 5x+1}{h} = \frac{5h}{h} = 5 \end{aligned}$$
- 79.** 
$$\begin{aligned} \frac{f(x+h) - f(x)}{h} &= \frac{[-7(x+h)+8] - [-7x+8]}{h} = \frac{[-7x-7h+8] - [-7x+8]}{h} \\ &= \frac{-7x-7h+8 + 7x-8}{h} = \frac{-7h}{h} = -7 \end{aligned}$$
- 80.** 
$$\begin{aligned} \frac{f(x+h) - f(x)}{h} &= \frac{[-8(x+h)-1] - [-8x-1]}{h} = \frac{[-8x-8h-1] - [-8x-1]}{h} \\ &= \frac{-8x-8h-1 + 8x+1}{h} = \frac{-8h}{h} = -8 \end{aligned}$$
- 81.** 
$$\begin{aligned} \frac{f(x+h) - f(x)}{h} &= \frac{[(x+h)^2 + 1] - [x^2 + 1]}{h} = \frac{[x^2 + 2xh + h^2 + 1] - [x^2 + 1]}{h} \\ &= \frac{x^2 + 2xh + h^2 + 1 - x^2 - 1}{h} \\ &= \frac{2xh + h^2}{h} = \frac{h(2x + h)}{h} = 2x + h \end{aligned}$$
- 82.** 
$$\begin{aligned} \frac{f(x+h) - f(x)}{h} &= \frac{[(x+h)^2 - 3] - [x^2 - 3]}{h} = \frac{[x^2 + 2xh + h^2 - 3] - [x^2 - 3]}{h} \\ &= \frac{x^2 + 2xh + h^2 - 3 - x^2 + 3}{h} \\ &= \frac{2xh + h^2}{h} = \frac{h(2x + h)}{h} = 2x + h \end{aligned}$$
- 83.** 
$$\begin{aligned} \frac{f(x+h) - f(x)}{h} &= \frac{[4(x+h)^2 - 6] - [4x^2 - 6]}{h} = \frac{[4x^2 + 8xh + 4h^2 - 6] - [4x^2 - 6]}{h} \\ &= \frac{4x^2 + 8xh + 4h^2 - 6 - 4x^2 + 6}{h} \\ &= \frac{8xh + 4h^2}{h} = \frac{h(8x + 4h)}{h} = 8x + 4h \end{aligned}$$

**EXERCISES 2.1**

$$\begin{aligned}
 84. \quad \frac{f(x+h) - f(x)}{h} &= \frac{[5(x+h)^2 + 3] - [5x^2 + 3]}{h} = \frac{[5x^2 + 10xh + 5h^2 + 3] - [5x^2 + 3]}{h} \\
 &= \frac{5x^2 + 10xh + 5h^2 + 3 - 5x^2 - 3}{h} \\
 &= \frac{10xh + 5h^2}{h} = \frac{h(10x + 5h)}{h} = 10x + 5h
 \end{aligned}$$

$$\begin{aligned}
 85. \quad \frac{f(x+h) - f(x)}{h} &= \frac{[(x+h)^2 + 3(x+h) - 7] - [x^2 + 3x - 7]}{h} \\
 &= \frac{[x^2 + 2xh + h^2 + 3x + 3h - 7] - [x^2 + 3x - 7]}{h} \\
 &= \frac{x^2 + 2xh + h^2 + 3x + 3h - 7 - x^2 - 3x + 7}{h} \\
 &= \frac{2xh + h^2 + 3h}{h} = \frac{h(2x + h + 3)}{h} = 2x + h + 3
 \end{aligned}$$

$$\begin{aligned}
 86. \quad \frac{f(x+h) - f(x)}{h} &= \frac{[(x+h)^2 - 5(x+h) + 1] - [x^2 - 5x + 1]}{h} \\
 &= \frac{[x^2 + 2xh + h^2 - 5x - 5h + 1] - [x^2 - 5x + 1]}{h} \\
 &= \frac{x^2 + 2xh + h^2 - 5x - 5h + 1 - x^2 + 5x - 1}{h} \\
 &= \frac{2xh + h^2 - 5h}{h} = \frac{h(2x + h - 5)}{h} = 2x + h - 5
 \end{aligned}$$

$$\begin{aligned}
 87. \quad \frac{f(x+h) - f(x)}{h} &= \frac{[2(x+h)^2 - 4(x+h) + 2] - [2x^2 - 4x + 2]}{h} \\
 &= \frac{[2x^2 + 4xh + 2h^2 - 4x - 4h + 2] - [2x^2 - 4x + 2]}{h} \\
 &= \frac{2x^2 + 4xh + 2h^2 - 4x - 4h + 2 - 2x^2 + 4x - 2}{h} \\
 &= \frac{4xh + 2h^2 - 4h}{h} = \frac{h(4x + 2h - 4)}{h} = 4x + 2h - 4
 \end{aligned}$$

$$\begin{aligned}
 88. \quad \frac{f(x+h) - f(x)}{h} &= \frac{[3(x+h)^2 + 2(x+h) - 3] - [3x^2 + 2x - 3]}{h} \\
 &= \frac{[3x^2 + 6xh + 3h^2 + 2x + 2h - 3] - [3x^2 + 2x - 3]}{h} \\
 &= \frac{3x^2 + 6xh + 3h^2 + 2x + 2h - 3 - 3x^2 - 2x + 3}{h} \\
 &= \frac{6xh + 3h^2 + 2h}{h} = \frac{h(6x + 3h + 2)}{h} = 6x + 3h + 2
 \end{aligned}$$

### EXERCISES 2.1

$$\begin{aligned}
 89. \quad \frac{f(x+h) - f(x)}{h} &= \frac{[-(x+h)^2 + (x+h) - 3] - [-x^2 + x - 3]}{h} \\
 &= \frac{[-x^2 - 2xh - h^2 + x + h - 3] - [-x^2 + x - 3]}{h} \\
 &= \frac{-x^2 - 2xh - h^2 + x + h - 3 + x^2 - x + 3}{h} \\
 &= \frac{-2xh - h^2 + h}{h} = \frac{h(-2x - h + 1)}{h} = -2x - h + 1
 \end{aligned}$$

$$\begin{aligned}
 90. \quad \frac{f(x+h) - f(x)}{h} &= \frac{[-3(x+h)^2 + 5(x+h) - 1] - [-3x^2 + 5x - 1]}{h} \\
 &= \frac{[-3x^2 - 6xh - 3h^2 + 5x + 5h - 1] - [-3x^2 + 5x - 1]}{h} \\
 &= \frac{-3x^2 - 6xh - 3h^2 + 5x + 5h - 1 + 3x^2 - 5x + 1}{h} \\
 &= \frac{-6xh - 3h^2 + 5h}{h} = \frac{h(-6x - 3h + 5)}{h} = -6x - 3h + 5
 \end{aligned}$$

$$\begin{aligned}
 91. \quad \frac{f(x+h) - f(x)}{h} &= \frac{(x+h)^3 - x^3}{h} = \frac{[x^3 + 3x^2h + 3xh^2 + h^3] - [x^3]}{h} \\
 &= \frac{3x^2h + 3xh^2 + h^3}{h} \\
 &= \frac{h(3x^2 + 3xh + h^2)}{h} = 3x^2 + 3xh + h^2
 \end{aligned}$$

$$\begin{aligned}
 92. \quad \frac{f(x+h) - f(x)}{h} &= \frac{-(x+h)^3 - (-x^3)}{h} = \frac{[-x^3 - 3x^2h - 3xh^2 - h^3] + x^3}{h} \\
 &= \frac{-3x^2h - 3xh^2 - h^3}{h} \\
 &= \frac{h(-3x^2 - 3xh - h^2)}{h} = -3x^2 - 3xh - h^2
 \end{aligned}$$

$$\begin{aligned}
 93. \quad \frac{f(x+h) - f(x)}{h} &= \frac{\frac{1}{x+h} - \frac{1}{x}}{h} = \frac{\left(\frac{1}{x+h} - \frac{1}{x}\right) \cdot x(x+h)}{h \cdot x(x+h)} \\
 &= \frac{x - (x+h)}{xh(x+h)} = \frac{-h}{xh(x+h)} = -\frac{1}{x(x+h)}
 \end{aligned}$$

$$94. \quad \frac{f(x+h) - f(x)}{h} = \frac{\sqrt{x+h} - \sqrt{x}}{h} \qquad \qquad \qquad 95. \quad f(x) = -0.6x + 132 \\
 f(25) = -0.6(25) + 132 = 117$$

$$\begin{aligned}
 96. \quad F(C) &= \frac{9}{5}C + 32 \\
 F(0) &= \frac{9}{5}(0) + 32 \quad F(-40) = \frac{9}{5}(-40) + 32 \quad F(10) = \frac{9}{5}(10) + 32 \\
 &= 32 \qquad \qquad \qquad = -40 \qquad \qquad \qquad = 50
 \end{aligned}$$

### EXERCISES 2.1

**97.**  $s(t) = -16t^2 + 10t + 300$   
 $s(3) = -16(3)^2 + 10(3) + 300 = 186 \text{ ft}$

**98.**  $v(t) = -32t + 15$   
 $v(t) = 0$   
 $-32t + 15 = 0$   
 $t = \frac{15}{32} \text{ seconds}$

**99.**  $g(d) = 300d$   
 $g(365) = 300(365) = 109,500 \text{ gallons}$

**100.**  $V(r) = \frac{4}{3}\pi r^3$   
 $V(29.5) = \frac{4}{3}\pi\left(\frac{29.5}{2}\right)^3 \approx 13,442 \text{ cm}^3$

**101.** Let  $x$  = the length. Then  $x + 5$  = the width.  
 $A(x) = x(x + 5) = x^2 + 5x$

**102.**  $V(x) = x(3x)(4) = 12x^2$

**103.** a.  $C(x) = 8x + 75$   
 b.  $C(85) = 8(85) + 75 = \$755$

**104.**  $C(x) = 45x + 60$

**105.** a.  $C(x) = 0.07x + 9.99$   
 b.  $C(20) = 0.07(20) + 9.99 = \$11.39$

**106.** a.  $I(x) = 2.5x - 40$   
 b.  $I(175) = 2.5(175) - 40 = \$397.50$

**107-110. Answers may vary.**

**111.** They are different. 10 is in the domain of  $f(x)$ , but not in the domain of  $g(x)$ .

**112-113. Answers may vary.**

**114.** They are different. The domain of  $f(x)$  is the set of all real numbers, but 3 is not in the domain of  $g(x)$ .

**115.**  $\frac{f(x+h) - f(x)}{h} = \frac{5-5}{h} = \frac{0}{h} = 0.$

**116.** e

**117.** d

**118.** b

**119.** e

**120.** f

**121.** c

**122.** e

**123.** a

### Exercises 2.2 (page 216)

**1.** quadrants

**2.** origin

**3.** to the right

**4.** upward

**5.** first

**6.** second

**7.** linear

**8.**  $y$ -axis

**9.**  $x$ -intercept

**10.** vertical

**11.** horizontal

**12.**  $\sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$

**13.** midpoint

**14.**  $M\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}\right)$

**15.**  $A(2, 3)$

**16.**  $B(-3, 5)$

**17.**  $C(-2, -3)$

**18.**  $D(4, -5)$

## EXERCISES 2.2


19.  $E(0, 0)$

20.  $F(-4, 0)$

21.  $G(-5, -5)$

22.  $H(2, -2)$

23, 25, 27, 29.




23. QI

25. QIII

27. QI

29. + x-axis

24, 26, 28, 30.



24. QII


26. QI

28. QIV

30. + y-axis


31.  $y - 2x = 7$   
 $y = 2x + 7$

x	y
0	7
-2	3




32.  $y + 3 = -4x$   
 $y = -4x - 3$

x	y
0	-3
-1	1




33.  $y + 5x = 5$   
 $y = -5x + 5$

x	y
0	5
1	0



34.  $y - 3x = 6$   
 $y = 3x + 6$


x	y
0	6
-2	0



## EXERCISES 2.2


**35.**  $6x - 3y = 10$   
 $-3y = -6x + 10$   
 $y = 2x - \frac{10}{3}$

x	y
0	$-\frac{10}{3}$
2	$\frac{2}{3}$




**37.**  $3x = 6y - 1$   
 $-6y = -3x - 1$   
 $y = \frac{1}{2}x + \frac{1}{6}$

x	y
0	$\frac{1}{6}$
-2	$-\frac{5}{6}$




**39.**  $2(x + y + 1) = x + 2$   
 $2x + 2y + 2 = x + 2$   
 $2y = -x$   
 $y = -\frac{1}{2}x$

x	y
0	0
-2	1




**36.**  $4x + 8y - 1 = 0$   
 $8y = -4x + 1$   
 $y = -\frac{1}{2}x + \frac{1}{8}$

x	y
0	$\frac{1}{8}$
4	$-\frac{15}{8}$




**38.**  $2x + 1 = 4y$   
 $-4y = -2x - 1$   
 $y = \frac{1}{2}x + \frac{1}{4}$

x	y
0	$\frac{1}{4}$
-2	$-\frac{3}{4}$




**40.**  $5(x + 2) = 3y - x$   
 $5x + 10 = 3y - x$   
 $-3y = -6x - 10$   
 $y = 2x + \frac{10}{3}$

x	y
0	$\frac{10}{3}$
-2	$-\frac{2}{3}$




## EXERCISES 2.2


41.  $x + y = 5$      $x + y = 5$   
 $x + 0 = 5$      $0 + y = 5$   
 $x = 5$              $y = 5$   
 $(5, 0)$              $(0, 5)$




43.  $2x - y = 4$      $2x - y = 4$   
 $2x - 0 = 4$      $2(0) - y = 4$   
 $2x = 4$              $-y = 4$   
 $x = 2$                $y = -4$   
 $(2, 0)$                $(0, -4)$



42.  $x - y = 3$      $x - y = 3$   
 $x - 0 = 3$      $0 - y = 3$   
 $x = 3$              $-y = 3$   
 $(3, 0)$              $y = -3$   
 $(0, -3)$




44.  $3x + y = 9$      $3x + y = 9$   
 $3x + 0 = 9$      $3(0) + y = 9$   
 $3x = 9$              $y = 9$   
 $x = 3$                $(0, 9)$   
 $(3, 0)$




## EXERCISES 2.2


**45.**  $3x + 2y = 6$        $3x + 2y = 6$   
 $3x + 2(0) = 6$        $3(0) + 2y = 6$   
 $3x = 6$        $2y = 6$   
 $x = 2$        $y = 3$   
 $(2, 0)$        $(0, 3)$




**47.**  $4x - 5y = 20$        $4x - 5y = 20$   
 $4x - 5(0) = 20$        $4(0) - 5y = 20$   
 $4x = 20$        $-5y = 20$   
 $x = 5$        $y = -4$   
 $(5, 0)$        $(0, -4)$




**46.**  $2x - 3y = 6$        $2x - 3y = 6$   
 $2x - 3(0) = 6$        $2(0) - 3y = 6$   
 $2x = 6$        $-3y = 6$   
 $x = 3$        $y = -2$   
 $(3, 0)$        $(0, -2)$




**48.**  $3x - 5y = 15$        $3x - 5y = 15$   
 $3x - 5(0) = 15$        $3(0) - 5y = 15$   
 $3x = 15$        $-5y = 15$   
 $x = 5$        $y = -3$   
 $(5, 0)$        $(0, -3)$



**49.**  $y = 3$




**50.**  $x = -4$



**51.**  $3x + 5 = -1$


$$3x = -6 \Rightarrow x = -2$$



## EXERCISES 2.2

52.  $7y - 1 = 6$


$$7y = 7 \Rightarrow y = 1$$



53.  $3(y + 2) = y$

$$3y + 6 = y$$


$$2y = -6 \Rightarrow y = -3$$



54.  $4 + 3y = 3(x + y)$

$$4 + 3y = 3x + 3y$$


$$4 = 3x \Rightarrow x = \frac{4}{3}$$



55.  $3(y + 2x) = 6x + y$

$$3y + 6x = 6x + y$$


$$2y = 0 \Rightarrow y = 0$$




56.  $5(y - x) = x + 5y$

$$5y - 5x = x + 5y$$

$$0 = 6x \Rightarrow x = 0$$




57.  $y = 3.7x - 4.5$



*x-int:*  $x = 1.22$

58.  $y = \frac{3}{5}x + \frac{5}{4}$




*x-int:*  $x = -2.08$

59.  $1.5x - 3y = 7$

$$-3y = -1.5x + 7$$


$$y = 0.5x - \frac{7}{3}$$



*x-int:*  $x = 4.67$

60.  $0.3x + y = 7.5$

$$y = -0.3x + 7.5$$



*x-int:*  $x = 25.00$

**EXERCISES 2.2**

$$\begin{aligned}
 61. \quad d &= \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} \\
 &= \sqrt{(4 - 0)^2 + (-3 - 0)^2} \\
 &= \sqrt{4^2 + (-3)^2} \\
 &= \sqrt{16 + 9} = \sqrt{25} = 5
 \end{aligned}$$

$$\begin{aligned}
 62. \quad d &= \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} \\
 &= \sqrt{(-5 - 0)^2 + (12 - 0)^2} \\
 &= \sqrt{(-5)^2 + (12)^2} \\
 &= \sqrt{25 + 144} = \sqrt{169} = 13
 \end{aligned}$$

$$\begin{aligned}
 63. \quad d &= \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} \\
 &= \sqrt{(-3 - 0)^2 + (2 - 0)^2} \\
 &= \sqrt{(-3)^2 + (2)^2} \\
 &= \sqrt{9 + 4} = \sqrt{13}
 \end{aligned}$$

$$\begin{aligned}
 64. \quad d &= \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} \\
 &= \sqrt{(5 - 0)^2 + (0 - 0)^2} \\
 &= \sqrt{(5)^2 + (0)^2} \\
 &= \sqrt{25 + 0} = \sqrt{25} = 5
 \end{aligned}$$

$$\begin{aligned}
 65. \quad d &= \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} \\
 &= \sqrt{(1 - 0)^2 + (1 - 0)^2} \\
 &= \sqrt{(1)^2 + (1)^2} \\
 &= \sqrt{1 + 1} = \sqrt{2}
 \end{aligned}$$

$$\begin{aligned}
 66. \quad d &= \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} \\
 &= \sqrt{(6 - 0)^2 + (-8 - 0)^2} \\
 &= \sqrt{(6)^2 + (-8)^2} \\
 &= \sqrt{36 + 64} = \sqrt{100} = 10
 \end{aligned}$$

$$\begin{aligned}
 67. \quad d &= \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} \\
 &= \sqrt{(\sqrt{3} - 0)^2 + (1 - 0)^2} \\
 &= \sqrt{(\sqrt{3})^2 + (1)^2} \\
 &= \sqrt{3 + 1} = \sqrt{4} = 2
 \end{aligned}$$

$$\begin{aligned}
 68. \quad d &= \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} \\
 &= \sqrt{(\sqrt{7} - 0)^2 + (\sqrt{2} - 0)^2} \\
 &= \sqrt{(\sqrt{7})^2 + (\sqrt{2})^2} \\
 &= \sqrt{7 + 2} = \sqrt{9} = 3
 \end{aligned}$$

$$\begin{aligned}
 69. \quad d &= \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} \\
 &= \sqrt{(3 - 6)^2 + (7 - 3)^2} \\
 &= \sqrt{(-3)^2 + (4)^2} \\
 &= \sqrt{9 + 16} = \sqrt{25} = 5
 \end{aligned}$$

$$\begin{aligned}
 70. \quad d &= \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} \\
 &= \sqrt{(4 - 9)^2 + (9 - 21)^2} \\
 &= \sqrt{(-5)^2 + (-12)^2} \\
 &= \sqrt{25 + 144} = \sqrt{169} = 13
 \end{aligned}$$

$$\begin{aligned}
 71. \quad d &= \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} \\
 &= \sqrt{[4 - (-1)]^2 + [-6 - 6]^2} \\
 &= \sqrt{(5)^2 + (-12)^2} \\
 &= \sqrt{25 + 144} = \sqrt{169} = 13
 \end{aligned}$$

$$\begin{aligned}
 72. \quad d &= \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} \\
 &= \sqrt{[0 - 6]^2 + [5 - (-3)]^2} \\
 &= \sqrt{(-6)^2 + (8)^2} \\
 &= \sqrt{36 + 64} = \sqrt{100} = 10
 \end{aligned}$$

**EXERCISES 2.2**

$$\begin{aligned}
 73. \quad d &= \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} \\
 &= \sqrt{[-2 - (-6)]^2 + [-15 - (-21)]^2} \\
 &= \sqrt{(4)^2 + (6)^2} \\
 &= \sqrt{16 + 36} = \sqrt{52} = 2\sqrt{13}
 \end{aligned}$$

$$\begin{aligned}
 74. \quad d &= \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} \\
 &= \sqrt{[-7 - (-11)]^2 + [11 - 7]^2} \\
 &= \sqrt{(4)^2 + (4)^2} \\
 &= \sqrt{16 + 16} = \sqrt{32} = 4\sqrt{2}
 \end{aligned}$$

$$\begin{aligned}
 75. \quad d &= \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} \\
 &= \sqrt{[3 - (-5)]^2 + [-3 - 5]^2} \\
 &= \sqrt{(8)^2 + (-8)^2} \\
 &= \sqrt{64 + 64} = \sqrt{128} = 8\sqrt{2}
 \end{aligned}$$

$$\begin{aligned}
 76. \quad d &= \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} \\
 &= \sqrt{[6 - (-3)]^2 + [-3 - 2]^2} \\
 &= \sqrt{(9)^2 + (-5)^2} \\
 &= \sqrt{81 + 25} = \sqrt{106}
 \end{aligned}$$

$$\begin{aligned}
 77. \quad d &= \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} \\
 &= \sqrt{[\pi - \pi]^2 + [-2 - 5]^2} \\
 &= \sqrt{(0)^2 + (-7)^2} \\
 &= \sqrt{0 + 49} = \sqrt{49} = 7
 \end{aligned}$$

$$\begin{aligned}
 78. \quad d &= \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} \\
 &= \sqrt{\left[\sqrt{5} - 0\right]^2 + [0 - 2]^2} \\
 &= \sqrt{\left(\sqrt{5}\right)^2 + (-2)^2} \\
 &= \sqrt{5 + 4} = \sqrt{9} = 3
 \end{aligned}$$

$$79. \quad M\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}\right) = M\left(\frac{2 + 6}{2}, \frac{4 + 8}{2}\right) = M\left(\frac{8}{2}, \frac{12}{2}\right) = M(4, 6)$$

$$80. \quad M\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}\right) = M\left(\frac{3 + (-1)}{2}, \frac{-6 + (-6)}{2}\right) = M\left(\frac{2}{2}, \frac{-12}{2}\right) = M(1, -6)$$

$$81. \quad M\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}\right) = M\left(\frac{2 + (-2)}{2}, \frac{-5 + 7}{2}\right) = M\left(\frac{0}{2}, \frac{2}{2}\right) = M(0, 1)$$

$$82. \quad M\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}\right) = M\left(\frac{0 + (-10)}{2}, \frac{3 + (-13)}{2}\right) = M\left(\frac{-10}{2}, \frac{-10}{2}\right) = M(-5, -5)$$

$$83. \quad M\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}\right) = M\left(\frac{-8 + 6}{2}, \frac{5 + (-4)}{2}\right) = M\left(\frac{-2}{2}, \frac{1}{2}\right) = M\left(-1, \frac{1}{2}\right)$$

$$84. \quad M\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}\right) = M\left(\frac{3 + 2}{2}, \frac{-2 + (-3)}{2}\right) = M\left(\frac{5}{2}, \frac{-5}{2}\right) = M\left(\frac{5}{2}, -\frac{5}{2}\right)$$

$$85. \quad M\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}\right) = M\left(\frac{0 + \sqrt{5}}{2}, \frac{0 + \sqrt{5}}{2}\right) = M\left(\frac{\sqrt{5}}{2}, \frac{\sqrt{5}}{2}\right)$$

## EXERCISES 2.2

**86.**  $M\left(\frac{x_1+x_2}{2}, \frac{y_1+y_2}{2}\right) = M\left(\frac{\sqrt{3}+0}{2}, \frac{0+(-\sqrt{5})}{2}\right) = M\left(\frac{\sqrt{3}}{2}, \frac{-\sqrt{5}}{2}\right) = M\left(\frac{\sqrt{3}}{2}, -\frac{\sqrt{5}}{2}\right)$

**87.** Let  $Q$  have coordinates  $(x, y)$ :

$$\begin{aligned} M\left(\frac{x_1+x_2}{2}, \frac{y_1+y_2}{2}\right) &= (3, 5) \\ \frac{x_1+x_2}{2} &= 3 \quad \frac{y_1+y_2}{2} = 5 \\ \frac{1+x}{2} &= 3 \quad \frac{4+y}{2} = 5 \\ 1+x &= 6 \quad 4+y = 10 \\ x &= 5 \quad y = 6 \\ Q(5, 6) \end{aligned}$$

**88.** Let  $Q$  have coordinates  $(x, y)$ :

$$\begin{aligned} M\left(\frac{x_1+x_2}{2}, \frac{y_1+y_2}{2}\right) &= (-5, 6) \\ \frac{x_1+x_2}{2} &= -5 \quad \frac{y_1+y_2}{2} = 6 \\ \frac{2+x}{2} &= -5 \quad \frac{-7+y}{2} = 6 \\ 2+x &= -10 \quad -7+y = 12 \\ x &= -12 \quad y = 19 \\ Q(-12, 19) \end{aligned}$$

**89.** Let  $Q$  have coordinates  $(x, y)$ :

$$\begin{aligned} M\left(\frac{x_1+x_2}{2}, \frac{y_1+y_2}{2}\right) &= (5, 5) \\ \frac{x_1+x_2}{2} &= 5 \quad \frac{y_1+y_2}{2} = 5 \\ \frac{5+x}{2} &= 5 \quad \frac{-5+y}{2} = 5 \\ 5+x &= 10 \quad -5+y = 10 \\ x &= 5 \quad y = 15 \\ Q(5, 15) \end{aligned}$$

**90.** Let  $Q$  have coordinates  $(x, y)$ :

$$\begin{aligned} M\left(\frac{x_1+x_2}{2}, \frac{y_1+y_2}{2}\right) &= (0, 0) \\ \frac{x_1+x_2}{2} &= 0 \quad \frac{y_1+y_2}{2} = 0 \\ \frac{-7+x}{2} &= 0 \quad \frac{3+y}{2} = 0 \\ -7+x &= 0 \quad 3+y = 0 \\ x &= 7 \quad y = -3 \\ Q(7, -3) \end{aligned}$$

**91.** Let the points be identified as  $A(13, -2)$ ,  $B(9, -8)$  and  $C(5, -2)$ .

$$AB = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} = \sqrt{(13 - 9)^2 + (-2 - (-8))^2} = \sqrt{16 + 36} = \sqrt{52} = 2\sqrt{13}$$

$$BC = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} = \sqrt{(9 - 5)^2 + (-8 - (-2))^2} = \sqrt{16 + 36} = \sqrt{52} = 2\sqrt{13}$$

Since  $AB$  and  $BC$  have the same length, the triangle is isosceles.

**92.** Let the points be identified as  $A(-1, 2)$ ,  $B(3, 1)$  and  $C(4, 5)$ .

$$AB = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} = \sqrt{(-1 - 3)^2 + (2 - 1)^2} = \sqrt{16 + 1} = \sqrt{17}$$

$$BC = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} = \sqrt{(3 - 4)^2 + (1 - 5)^2} = \sqrt{1 + 16} = \sqrt{17}$$

Since  $AB$  and  $BC$  have the same length, the triangle is isosceles.

**93.**  $M = \left(\frac{2+6}{2}, \frac{4+10}{2}\right) = \left(\frac{8}{2}, \frac{14}{2}\right) = (4, 7); N = \left(\frac{4+6}{2}, \frac{6+10}{2}\right) = \left(\frac{10}{2}, \frac{16}{2}\right) = (5, 8)$

$$MN = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} = \sqrt{(4 - 5)^2 + (7 - 8)^2} = \sqrt{1 + 1} = \sqrt{2}$$

## EXERCISES 2.2

**94.**  $M = \left( \frac{0+b}{2}, \frac{0+c}{2} \right) = \left( \frac{b}{2}, \frac{c}{2} \right); N = \left( \frac{a+b}{2}, \frac{0+c}{2} \right) = \left( \frac{a+b}{2}, \frac{c}{2} \right)$   
 $AB = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} = \sqrt{(0-a)^2 + (0-0)^2} = \sqrt{a^2} = a$   
 $MN = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} = \sqrt{\left( \frac{b}{2} - \frac{a+b}{2} \right)^2 + \left( \frac{c}{2} - \frac{c}{2} \right)^2} = \sqrt{\frac{a^2}{4} + 0} = \frac{a}{2} = \frac{1}{2}AB$

**95.**  $M = \left( \frac{0+a}{2}, \frac{b+0}{2} \right) = \left( \frac{a}{2}, \frac{b}{2} \right); L = \left( \frac{a}{2}, 0 \right); N = \left( 0, \frac{b}{2} \right)$   
 Area of  $AOB = \frac{1}{2} \cdot \text{base} \cdot \text{height} = \frac{1}{2}(OA)(OB) = \frac{1}{2}(a)(b) = \frac{1}{2}ab$   
 Area of  $OLMN = \text{length} \cdot \text{width} = (OL)(ON) = \frac{a}{2} \cdot \frac{b}{2} = \frac{1}{4}ab = \frac{1}{2}(\text{Area of } AOB)$

- 96.** Let  $x =$  the width (from  $A$  to  $D$ ). Then  
 the length (from  $A$  to  $B$ ) =  $2x$ .

$$\text{Perimeter} = 42$$

$$x + 2x + x + 2x = 42$$

$$6x = 42$$

$$x = 7$$

Thus, the distance from  $A$  to  $D$  is 7 and the distance from  $A$  to  $B$  is  $2(7) = 14$ . Thus, the  $x$ -coordinate of  $C$  is  $-3 + 14$ , or 11. The  $y$ -coordinate of  $C$  is  $-2 + 7$ , or 5. Point  $C$  then has coordinates  $(11, 5)$ .

**97.**  $y = 17500x + 325000$

$$y = 17500(5) + 325000$$

$$y = 87500 + 325000$$

$$y = 412500$$

The value will be \$412,500.

**98.** Set  $y = 0$ :

$$y = -1920x + 24,000$$

$$0 = -1920x + 24,000$$

$$1920x = 24,000$$

$$x = 12.5$$

The car will be worthless after 12.5 years.

**99.**  $p = -\frac{1}{10}q + 170$

$$150 = -\frac{1}{10}q + 170$$

$$\frac{1}{10}q = 20$$

$$q = 200$$

200 scanners will be sold.

**100.**  $p = \frac{1}{10}q + 130$

$$150 = \frac{1}{10}q + 130$$

$$20 = \frac{1}{10}q$$

$$200 = q$$

200 TVs will be produced.

**101.**  $V = \frac{nv}{N}$

$$60 = \frac{12v}{20}$$

$$1200 = 12v$$

$$100 = v$$

The smaller gear is spinning at 100 rpm.

**102.**  $f(x) = 430 - 0.005x$

$$350 = 430 - 0.005d$$

$$0.005d = 80$$

$$d = 16,000$$

\$16,000 would reduce the number to 350.

## EXERCISES 2.2


**103.**  $d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$   
 $= \sqrt{[0 - 30]^2 + [10 - 25]^2}$   
 $= \sqrt{(-30)^2 + (-15)^2}$   
 $= \sqrt{900 + 225} = \sqrt{1125} = 15\sqrt{5}$  yd

**104.**  $d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$   
 $= \sqrt{[90 - 0]^2 + [90 - 0]^2}$   
 $= \sqrt{(90)^2 + (90)^2}$   
 $= \sqrt{8100 + 8100} = \sqrt{16200} = 90\sqrt{2}$  ft

**105.**  $M\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}\right) = M\left(\frac{30 + 0}{2}, \frac{25 + 10}{2}\right) = M\left(\frac{30}{2}, \frac{35}{2}\right) = M(15, 17.5)$


**106.**  $M\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}\right) = M\left(\frac{90 + 0}{2}, \frac{90 + 0}{2}\right) = M\left(\frac{90}{2}, \frac{90}{2}\right) = M(45, 45)$

**107.**



$$\begin{aligned} d^2 &= 70^2 + 156^2 \\ d^2 &= 4900 + 24,336 \\ d^2 &= 29,236 \\ d &= \sqrt{29,236} \\ d &\approx 171 \text{ miles} \end{aligned}$$

**108.**



$$\begin{aligned} d^2 &= 10^2 + 3^2 \\ d^2 &= 100 + 9 \\ d^2 &= 109 \\ d &= \sqrt{109} \\ d &\approx 10.4 \text{ mm} \end{aligned}$$

**109-112. Answers may vary.**

**113.** True.

**114.** True.

**115.** False. Vertical lines have equations that are not functions.

**116.** False. Only two points are required to graph a line.

**117.** False. The vertical line  $x = 0$  has infinitely many  $y$ -intercepts.

**118.** False. Most horizontal lines have no  $x$ -intercept.

**119.** True.

**120.** True.

## Exercises 2.3 (page 229)

**1.** divided

**2.**  $y$

**3.** run

**4.** same order

**5.** the change in

**6.** horizontal

**7.** vertical

**8.** parallel

**9.** perpendicular

**10.**  $-1$

**11.**  $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{2 - (-1)}{2 - (-1)} = \frac{3}{3} = 1$

**12.**  $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{3 - (-1)}{5 - 3} = \frac{4}{2} = 2$

**13.**  $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{-2 - 3}{6 - (-6)} = \frac{-5}{12} = -\frac{5}{12}$

**14.**  $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{10 - 5}{3 - 2} = \frac{5}{1} = 5$

### EXERCISES 2.3

15.  $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{5 - (-2)}{-1 - 3} = \frac{7}{-4} = -\frac{7}{4}$

16.  $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{16 - 7}{6 - 3} = \frac{9}{3} = 3$

17.  $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{1 - (-7)}{4 - 8} = \frac{8}{-4} = -2$

18.  $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{17 - 17}{17 - 5} = \frac{0}{12} = 0$

19.  $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{-14 - (-14)}{2 - (-7)} = \frac{0}{9} = 0$

20.  $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{-3 - 3}{-4 - (-4)} = \frac{-6}{0} \Rightarrow \text{und.}$

21.  $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{-2 - 3}{-5 - (-5)} = \frac{-5}{0} \Rightarrow \text{und.}$

22.  $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{2 - \sqrt{7}}{\sqrt{7} - 2} = -1$

23.  $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{\frac{7}{3} - \frac{2}{3}}{\frac{5}{2} - \frac{3}{2}} = \frac{\frac{5}{3}}{\frac{2}{2}} = \frac{\frac{5}{3}}{1} = \frac{5}{3}$

24.  $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{-\frac{5}{3} - \frac{1}{3}}{\frac{3}{5} - \left(-\frac{2}{5}\right)} = \frac{-\frac{6}{3}}{\frac{5}{5}} = -2$

25.  $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{a - c}{(b + c) - (a + b)}$   
 $= \frac{a - c}{c - a} = -1$

26.  $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{a - 0}{(a + b) - b} = \frac{a}{a} = 1$

27.  $y = 3x + 2 \quad m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{5 - 2}{1 - 0}$   
 $\begin{array}{|c|c|} \hline x & y \\ \hline 0 & 2 \\ \hline 1 & 5 \\ \hline \end{array} \quad = \frac{3}{1} = 3$

28.  $y = 5x - 8 \quad m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{-3 - (-8)}{1 - 0}$   
 $\begin{array}{|c|c|} \hline x & y \\ \hline 0 & -8 \\ \hline 1 & -3 \\ \hline \end{array} \quad = \frac{5}{1} = 5$

29.  $y = 4x - 6 \quad m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{-2 - (-6)}{1 - 0}$   
 $\begin{array}{|c|c|} \hline x & y \\ \hline 0 & -6 \\ \hline 1 & -2 \\ \hline \end{array} \quad = \frac{4}{1} = 4$

30.  $y = -\frac{1}{3}x + 5 \quad m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{4 - 5}{3 - 0}$   
 $\begin{array}{|c|c|} \hline x & y \\ \hline 0 & 5 \\ \hline 3 & 4 \\ \hline \end{array} \quad = \frac{-1}{3} = -\frac{1}{3}$

31.  $5x - 10y = 3 \quad m = \frac{y_2 - y_1}{x_2 - x_1}$   
 $\begin{array}{|c|c|} \hline x & y \\ \hline 0 & -\frac{3}{10} \\ \hline 1 & \frac{1}{5} \\ \hline \end{array} \quad = \frac{\frac{1}{5} - \left(-\frac{3}{10}\right)}{1 - 0}$   
 $= \frac{\frac{5}{10}}{1} = \frac{1}{2}$

32.  $8y + 2x = 5 \quad m = \frac{y_2 - y_1}{x_2 - x_1}$   
 $\begin{array}{|c|c|} \hline x & y \\ \hline 0 & \frac{5}{8} \\ \hline 1 & \frac{3}{8} \\ \hline \end{array} \quad = \frac{\frac{3}{8} - \frac{5}{8}}{1 - 0}$   
 $= \frac{-\frac{2}{8}}{1} = -\frac{1}{4}$

33.  $3(y + 2) = 2x - 3 \quad m = \frac{y_2 - y_1}{x_2 - x_1}$   
 $3y - 2x = -9 \quad = \frac{-1 - (-3)}{3 - 0}$   
 $\begin{array}{|c|c|} \hline x & y \\ \hline 0 & -3 \\ \hline 3 & -1 \\ \hline \end{array} \quad = \frac{2}{3}$

34.  $4(x - 2) = 3y + 2 \quad m = \frac{y_2 - y_1}{x_2 - x_1}$   
 $4x - 3y = 10 \quad = \frac{-2 - \left(-\frac{10}{3}\right)}{1 - 0}$   
 $\begin{array}{|c|c|} \hline x & y \\ \hline 0 & -\frac{10}{3} \\ \hline 1 & -2 \\ \hline \end{array} \quad = \frac{\frac{4}{3}}{1} = \frac{4}{3}$

### EXERCISES 2.3

**35.**  $3(y + x) = 3(x - 1)$      $m = \frac{y_2 - y_1}{x_2 - x_1}$

$$\begin{aligned} 3y &= -3 \\ y &= -1 \end{aligned}$$

$x$	$y$
0	-1
1	-1

$$\begin{aligned} &= \frac{-1 - (-1)}{1 - 0} \\ &= \frac{0}{1} = 0 \end{aligned}$$

**36.**  $2x + 5 = 2(y + x)$      $m = \frac{y_2 - y_1}{x_2 - x_1}$

$$\begin{aligned} 5 &= 2y \\ \frac{5}{2} &= y \end{aligned}$$

$x$	$y$
0	$\frac{5}{2}$
1	$\frac{5}{2}$

$$\begin{aligned} &= \frac{\frac{5}{2} - \frac{5}{2}}{1 - 0} \\ &= \frac{0}{1} = 0 \end{aligned}$$

**37.** horizontal  $\Rightarrow m = 0$

**38.**  $2y = 5$

$$y = \frac{5}{2}$$

horizontal  $\Rightarrow m = 0$

**39.**  $f(x) = \frac{1}{4} \Rightarrow y = \frac{1}{4}$

horizontal  $\Rightarrow m = 0$

**40.**  $f(x) = \pi \Rightarrow y = \pi$   
horizontal  $\Rightarrow m = 0$

**41.**  $x = -\frac{1}{2}$   
vertical  $\Rightarrow m$  is undefined.

**42.**  $x - 7 = 0$   
 $x = 7$   
vertical  $\Rightarrow m$  is undefined.

**43.** The slope is negative.

**44.** The slope is zero.

**45.** The slope is positive.

**46.** The slope is positive.

**47.** The slope is undefined.

**48.** The slope is negative.

**49.**  $m_1 m_2 = 3\left(-\frac{1}{3}\right) = -1$   
perpendicular

**50.**  $m_1 \neq m_2$ ;  $m_1 m_2 = \frac{2}{3} \cdot \frac{3}{2} = 1 \neq -1$   
neither

**51.**  $m_1 = \sqrt{8} = 2\sqrt{2} = m_2$   
parallel

**52.**  $m_1 m_2 = 1(-1) = -1$   
perpendicular

**53.**  $m_1 m_2 = -\sqrt{2}\left(\frac{\sqrt{2}}{2}\right) = -1$   
perpendicular

**54.**  $m_2 = \sqrt{28} = 2\sqrt{7} = m_1$   
parallel

**55.**  $m_1 m_2 = -0.125(8) = -1$   
perpendicular

**56.**  $m_1 = 0.125 = \frac{1}{8} = m_2$   
parallel

**57.**  $m_1 m_2 = ab^{-1}(-a^{-1}b) = -a^0 b^0 = -1$   
perpendicular

**58.**  $m_1 = \left(\frac{a}{b}\right)^{-1} = \frac{b}{a} \neq -\frac{b}{a} = m_2$   
 $m_1 m_2 = \frac{b}{a} \left(-\frac{b}{a}\right) \neq -1 \Rightarrow \text{neither}$

### EXERCISES 2.3

For Exercises 59-64 use the slope of line through R and S calculated below:

$$m_{RS} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{7 - 5}{2 - (-3)} = \frac{2}{5}$$

**59.**  $m_{PQ} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{6 - 4}{7 - 2} = \frac{2}{5} = m_{RS} \Rightarrow$  parallel

**60.**  $m_{PQ} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{4 - 8}{-13 - (-3)} = \frac{-4}{-10} = \frac{2}{5} = m_{RS} \Rightarrow$  parallel

**61.**  $m_{PQ} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{1 - 6}{-2 - (-4)} = \frac{-5}{2} = -\frac{5}{2} \Rightarrow$  perpendicular

**62.**  $m_{PQ} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{1 - (-9)}{4 - 0} = \frac{10}{4} = \frac{5}{2} \Rightarrow$  neither

**63.**  $m_{PQ} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{6a - a}{3a - a} = \frac{5a}{2a} = \frac{5}{2} \Rightarrow$  neither

**64.**  $m_{PQ} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{6b - b}{-b - b} = \frac{5b}{-2b} = -\frac{5}{2} \Rightarrow$  perpendicular

**65.**  $m_{PQ} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{9 - 7}{2 - (-3)} = \frac{2}{5}; m_{RS} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{-6 - (-4)}{x - 10} = \frac{-2}{x - 10}$   
 $\frac{2}{5} \cdot \frac{-1}{-1} = \frac{-2}{-5}; x - 10 = -5 \Rightarrow \boxed{x = 5}$

**66.**  $m_{PQ} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{7 - (-3)}{5 - 2} = \frac{10}{3}; m_{RS} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{y - (-1)}{6 - 3} = \frac{y + 1}{3}$   
 $10 = y + 1 \Rightarrow \boxed{y = 9}$

**67.**  $m_{PQ} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{0 - (-7)}{1 - 2} = \frac{7}{-1} = -7; m_{RS} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{y - 5}{-2 - (-9)} = \frac{y - 5}{7}$   
 $-7 = \frac{-7}{1}; \text{Perp. slope} = \frac{1}{7}; y - 5 = 1 \Rightarrow \boxed{y = 6}$

**68.**  $m_{PQ} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{4 - (-2)}{3 - 1} = \frac{6}{2} = 3; m_{RS} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{5 - 6}{6 - x} = \frac{-1}{6 - x}$   
 $3 = \frac{3}{1}; \text{Perp. slope} = -\frac{1}{3} = \frac{-1}{3}; 6 - x = 3 \Rightarrow \boxed{x = 3}$

**69.**  $m_{PQ} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{9 - 8}{-6 - (-2)} = \frac{1}{-4} = -\frac{1}{4}$   
 $m_{PR} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{5 - 8}{2 - (-2)} = \frac{-3}{4} = -\frac{3}{4} \Rightarrow$  not on same line

### EXERCISES 2.3

**70.**  $m_{PQ} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{-2 - (-1)}{3 - 1} = \frac{-1}{2} = -\frac{1}{2}$   
 $m_{PR} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{0 - (-1)}{-3 - 1} = \frac{1}{-4} = -\frac{1}{4} \Rightarrow$  not on same line

**71.**  $m_{PQ} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{0 - a}{0 - (-a)} = \frac{-a}{a} = -1$   
 $m_{PR} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{-a - a}{a - (-a)} = \frac{-2a}{2a} = -1 \Rightarrow$  on same line

**72.**  $m_{PQ} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{b - (a + b)}{(a + b) - a} = \frac{-a}{b} = -\frac{a}{b}$   
 $m_{PR} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{a - (a + b)}{(a - b) - a} = \frac{-b}{-b} = 1 \Rightarrow$  not on same line

**73.**  $m_{PQ} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{-5 - 4}{2 - 5} = \frac{-9}{-3} = 3$   
 $m_{PR} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{-3 - 4}{8 - 5} = \frac{-7}{3} = -\frac{7}{3}$   
 $m_{QR} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{-3 - (-5)}{8 - 2} = \frac{2}{6} = \frac{1}{3} \Rightarrow$  None are perpendicular.

**74.**  $m_{PQ} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{6 - (-2)}{4 - 8} = \frac{8}{-4} = -2$   
 $m_{PR} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{7 - (-2)}{6 - 8} = \frac{9}{-2} = -\frac{9}{2}$   
 $m_{QR} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{7 - 6}{6 - 4} = \frac{1}{2} \Rightarrow PQ$  and  $QR$  are perpendicular.

**75.**  $m_{PQ} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{9 - 3}{1 - 1} = \frac{6}{0} \Rightarrow$  undefined  $\Rightarrow$  vertical  
 $m_{PR} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{3 - 3}{7 - 1} = \frac{0}{6} = 0 \Rightarrow$  horizontal  
 $m_{QR} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{3 - 9}{7 - 1} = \frac{-6}{6} = -1 \Rightarrow PQ$  and  $PR$  are perpendicular.

**76.**  $m_{PQ} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{2 - (-3)}{-3 - 2} = \frac{5}{-5} = -1$   
 $m_{PR} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{8 - (-3)}{3 - 2} = \frac{11}{1} = 11$   
 $m_{QR} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{8 - 2}{3 - (-3)} = \frac{6}{6} = 1 \Rightarrow PQ$  and  $QR$  are perpendicular.

### EXERCISES 2.3

77.  $m_{PQ} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{b - 0}{a - 0} = \frac{b}{a}$

$$m_{PR} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{a - 0}{-b - 0} = \frac{a}{-b} = -\frac{a}{b}$$

$m_{QR} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{a - b}{-b - a} = \frac{a - b}{-b - a} \Rightarrow PQ \text{ and } PR \text{ are perpendicular.}$

78.  $m_{PQ} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{a - b}{-b - a} = \frac{a - b}{-(a + b)} = -\frac{a - b}{a + b}$

$$m_{PR} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{(a + b) - b}{(a - b) - a} = \frac{a}{-b} = -\frac{a}{b}$$

$m_{QR} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{(a + b) - a}{(a - b) - (-b)} = \frac{b}{a} \Rightarrow PR \text{ and } QR \text{ are perpendicular.}$

79.  $m_{AB} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{4 - (-1)}{-3 - (-1)} = \frac{5}{-2} = -\frac{5}{2}$

$m_{AC} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{1 - (-1)}{4 - (-1)} = \frac{2}{5} \Rightarrow AB \text{ and } AC \text{ are perpendicular.} \Rightarrow \text{right triangle}$

80.  $m_{DE} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{3 - 1}{-1 - 0} = \frac{2}{-1} = -2$

$m_{EF} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{5 - 3}{3 - (-1)} = \frac{2}{4} = \frac{1}{2} \Rightarrow DE \text{ and } EF \text{ are perpendicular.} \Rightarrow \text{right triangle}$

81.  $m_{AB} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{0 - (-1)}{3 - 1} = \frac{1}{2}; \quad d(A, B) = \sqrt{(1 - 3)^2 + (-1 - 0)^2} = \sqrt{5}$

$$m_{BC} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{2 - 0}{2 - 3} = \frac{2}{-1} = -2; \quad d(B, C) = \sqrt{(3 - 2)^2 + (0 - 2)^2} = \sqrt{5}$$

$$m_{CD} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{1 - 2}{0 - 2} = \frac{-1}{-2} = \frac{1}{2}; \quad d(C, D) = \sqrt{(2 - 0)^2 + (2 - 1)^2} = \sqrt{5}$$

$$m_{DA} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{1 - (-1)}{0 - 1} = \frac{2}{-1} = -2; \quad d(D, A) = \sqrt{(1 - 0)^2 + (-1 - 1)^2} = \sqrt{5}$$

Adjacent sides are perpendicular and congruent, so the figure is a square.

82.  $m_{EF} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{0 - (-1)}{3 - (-1)} = \frac{1}{4}; \quad d(E, F) = \sqrt{(-1 - 3)^2 + (-1 - 0)^2} = \sqrt{17}$

$$m_{FG} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{4 - 0}{2 - 3} = \frac{4}{-1} = -4; \quad d(F, G) = \sqrt{(3 - 2)^2 + (0 - 4)^2} = \sqrt{17}$$

$$m_{GH} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{3 - 4}{-2 - 2} = \frac{-1}{-4} = \frac{1}{4}; \quad d(G, H) = \sqrt{(2 - (-2))^2 + (4 - 3)^2} = \sqrt{17}$$

$$m_{HE} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{3 - (-1)}{-2 - (-1)} = \frac{4}{-1} = -4; \quad d(H, E) = \sqrt{(-1 - (-2))^2 + (-1 - 3)^2} = \sqrt{17}$$

Adjacent sides are perpendicular and congruent, so the figure is a square.

### EXERCISES 2.3

**83.**  $m_{AB} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{3 - (-2)}{3 - (-2)} = \frac{5}{5} = 1$

$$m_{BC} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{6 - 3}{2 - 3} = \frac{3}{-1} = -3$$

$$m_{CD} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{1 - 6}{-3 - 2} = \frac{-5}{-5} = 1$$

$$m_{DA} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{1 - (-2)}{-3 - (-2)} = \frac{3}{-1} = -3$$

Opposite sides are parallel, so the figure is a parallelogram.

**84.**  $m_{EF} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{1 - (-2)}{5 - 1} = \frac{3}{4}$

$$m_{FG} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{4 - 1}{3 - 5} = \frac{3}{-2} = -\frac{3}{2}$$

$$m_{GH} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{4 - 4}{-3 - 3} = \frac{0}{-6} = 0$$

$$m_{HE} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{4 - (-2)}{-3 - 1} = \frac{6}{-4} = -\frac{3}{2}$$

Exactly one pair of sides is parallel, so the figure is a trapezoid.

**85.**  $M\left(\frac{5+7}{2}, \frac{9+5}{2}\right) = M\left(\frac{12}{2}, \frac{14}{2}\right) = M(6, 7); N\left(\frac{1+7}{2}, \frac{3+5}{2}\right) = N\left(\frac{8}{2}, \frac{8}{2}\right) = N(4, 4)$

$$m_{MN} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{4 - 7}{4 - 6} = \frac{-3}{-2} = \frac{3}{2}; m_{AC} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{9 - 3}{5 - 1} = \frac{6}{4} = \frac{3}{2} \Rightarrow MN \parallel AC$$

**86.**  $d(AB) = \sqrt{(0-a)^2 + (0-0)^2} = \sqrt{a^2 + 0^2} = \sqrt{a^2} = a$

$$d(AC) = \sqrt{(0-b)^2 + (0-c)^2} = \sqrt{b^2 + c^2}. \text{ From the given information, } a = \sqrt{b^2 + c^2}.$$

$$m_{AD} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{c - 0}{(a+b) - 0} = \frac{c}{a+b}; m_{BC} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{0 - c}{a - b} = \frac{-c}{a - b}$$

$$m_{AD}m_{BC} = \frac{c}{a+b} \cdot \frac{-c}{a-b} = \frac{-c^2}{a^2 - b^2} = \frac{-c^2}{(\sqrt{b^2 + c^2})^2 - b^2} = \frac{-c^2}{b^2 + c^2 - b^2} = \frac{-c^2}{c^2} = -1$$

Thus,  $AD$  is perpendicular to  $BC$ .

**87.**  $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{42 - 14}{5 - 1} = \frac{28}{4} = 7$

The rate of growth was 7 students per year.

**88.**  $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{110,000 - 50,000}{3 - 1}$

$$= \frac{60,000}{2} = 30,000$$

The sales increased \$30,000 per year.

**89.**  $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{6700 - 2200}{10 - 3}$   
 $= \frac{4500}{7} \approx 642.86$

The cost decreased about \$642.86 per year.


**90.** The cost absorbed by the hospital was \$247 in 2000, \$375 in 2005 and \$505 in 2010.

$$m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{505 - 247}{2010 - 2000} = \frac{258}{10} = 25.8$$


The cost absorbed by the hospital increased by \$25.80 per year.

### EXERCISES 2.3


91.  $\frac{\Delta T}{\Delta t} = \frac{T_2 - T_1}{t_2 - t_1}$  = the hourly rate of change of temperature.  
(Let  $t = x$  and  $T = y$ .)




92.  $\frac{\Delta D}{\Delta d} = \frac{D_2 - D_1}{d_2 - d_1}$  = the daily rate of change of the Dow Jones average.



93.  $D = 590t$ ; The slope is the speed of the plane.



94.  $A = 25n$ ; The slope is the monthly increase of the value of the account.



95-98. Answers may vary.

99. False.  $m = \frac{y_2 - y_1}{x_2 - x_1}$ .

100. True.

101. True. ( $\Delta y = 0$ .)

102. True. ( $\Delta x = 0$ .)

103. False. The line will be horizontal, so the slope is 0.

104. False. The line will be vertical, so the slope is undefined.

105.  $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{10.25 - 6.95}{2014 - 2008} = \frac{3.30}{6} = 0.55$ . True.

106.  $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{9139 - 6015}{2014 - 2010} = \frac{3124}{4} = 781$ . True.

**EXERCISES 2.4**

**Exercises 2.4** (page 243)

1. slope-intercept

2.  $m$

3.  $y$ -intercept

4.  $y - y_1 = m(x - x_1)$

5.  $Ax + By = C$

6. regression

7.  $y = mx + b$

$y = 3x - 2$

8.  $y = mx + b$

$y = -\frac{1}{3}x + \frac{2}{3}$

9.  $y = mx + b$

$y = 5x - \frac{1}{5}$

10.  $y = mx + b$

$y = \sqrt{2}x + \sqrt{2}$

11.  $y = mx + b$

$y = ax + \frac{1}{a}$

12.  $y = mx + b$

$y = ax + 2a$

13.  $y = mx + b$

$y = ax + a$

14.  $y = mx + b$

$y = \frac{1}{a}x + a$

15.  $y = mx + b$

$0 = \frac{3}{2}(0) + b$

$0 = b$

$y = mx + b$

$y = \frac{3}{2}x + 0$

$2y = 3x$

$-3x + 2y = 0$

$3x - 2y = 0$

16.  $y = mx + b$

$-7 = -\frac{2}{3}(-3) + b$

$-7 = 2 + b$

$-9 = b$

$y = mx + b$

$y = -\frac{2}{3}x - 9$

$3y = -2x - 27$

$2x + 3y = -27$

17.  $y = mx + b$

$5 = -3(-3) + b$

$5 = 9 + b$

$-4 = b$

$y = mx + b$

$y = -3x - 4$

$3x + y = -4$

18.  $y = mx + b$

$1 = 1(-5) + b$

$1 = -5 + b$

$6 = b$

$y = mx + b$

$y = 1x + 6$

$-x + y = 6$

$x - y = -6$

19.  $y = mx + b$

$\sqrt{2} = \sqrt{2}(0) + b$

$\sqrt{2} = b$

$y = mx + b$

$y = \sqrt{2}x + \sqrt{2}$

$-\sqrt{2}x + y = \sqrt{2}$

$\sqrt{2}x - y = -\sqrt{2}$

20.  $y = mx + b$

$0 = 2\sqrt{3}(-\sqrt{3}) + b$

$0 = -6 + b$

$6 = b$

$y = mx + b$

$y = 2\sqrt{3}x + 6$

$-2\sqrt{3}x + y = 6$

$2\sqrt{3}x - y = -6$

21.  $3x - 2y = 8$

$-2y = -3x + 8$

$y = \frac{3}{2}x - 4$

$m = \frac{3}{2}, (0, -4)$

22.  $-2x + 4y = 12$

$4y = 2x + 12$

$y = \frac{1}{2}x + 3$

$m = \frac{1}{2}, (0, 3)$

23.  $-2(x + 3y) = 5$

$-2x - 6y = 5$

$-6y = 2x + 5$

$y = -\frac{1}{3}x - \frac{5}{6}$

$m = -\frac{1}{3}, \left(0, -\frac{5}{6}\right)$

### EXERCISES 2.4

24.  $5(2x - 3y) = 4$

$$10x - 15y = 4$$

$$-15y = -10x + 4$$

$$y = \frac{2}{3}x - \frac{4}{15}$$

$$m = \frac{2}{3}, \left(0, -\frac{4}{15}\right)$$

25.  $x = \frac{2y - 4}{7}$

$$7x = 2y - 4$$

$$-2y = -7x - 4$$

$$y = \frac{7}{2}x + 2$$

$$m = \frac{7}{2}, (0, 2)$$

26.  $3x + 4 = -\frac{2(y - 3)}{5}$

$$15x + 20 = -2(y - 3)$$

$$15x + 20 = -2y + 6$$


$$2y = -15x - 14$$

$$y = -\frac{15}{2}x - 7$$

$$m = -\frac{15}{2}, (0, -7)$$


27.  $x - y = 1$

$$y = x - 1 \Rightarrow m = 1, (0, -1)$$



28.  $x + y = 2$

$$y = -x + 2 \Rightarrow m = -1, (0, 2)$$




29.  $x = \frac{3}{2}y - 3$

$$2x = 3y - 6$$

$$-3y = -2x - 6$$

$$y = \frac{2}{3}x + 2 \Rightarrow m = \frac{2}{3}, (0, 2)$$




30.  $x = -\frac{4}{5}y + 2$

$$5x = -4y + 10$$

$$4y = -5x + 10$$

$$y = -\frac{5}{4}x + \frac{5}{2} \Rightarrow m = -\frac{5}{4}, \left(0, \frac{5}{2}\right)$$




### EXERCISES 2.4

31.  $3(y - 4) = -2(x - 3)$

$$3y - 12 = -2x + 6$$

$$3y = -2x + 18$$

$$y = -\frac{2}{3}x + 6 \Rightarrow m = -\frac{2}{3}, (0, 6)$$




32.  $-4(2x + 3) = 3(3y + 8)$

$$-8x - 12 = 9y + 24$$

$$-9y = 8x + 36$$

$$y = -\frac{8}{9}x - 4: m = -\frac{8}{9}, (0, -4)$$



33.  $y = 3x + 4 \quad y = 3x - 7$

$$m = 3 \quad m = 3$$

The lines are parallel.

35.  $x + y = 2 \quad y = x + 5$   
 $y = -x + 2 \quad m = 1$

$$m = -1$$

The lines are perpendicular.

34.  $y = 4x - 13 \quad y = \frac{1}{4}x + 13$

$$m = 4$$

$$m = \frac{1}{4}$$

The lines are neither.

36.  $x = y + 2 \quad y = x + 3$   
 $-y = -x + 2 \quad m = 1$   
 $y = x - 2$

$$m = 1$$

The lines are parallel.

37.  $y = 3x + 7 \quad 2y = 6x - 9$   
 $m = 3 \quad y = 3x - \frac{9}{2}$   
 $m = 3$

The lines are parallel.

38.  $2x + 3y = 9 \quad 3x - 2y = 5$   
 $3y = -2x + 9 \quad -2y = -3x + 5$   
 $y = -\frac{2}{3}x + 3 \quad y = \frac{3}{2}x - \frac{5}{2}$

$$m = -\frac{2}{3} \quad m = \frac{3}{2}$$

The lines are perpendicular.

### EXERCISES 2.4

**39.**  $3x + 6y = 1$        $y = \frac{1}{2}x - \frac{1}{6}$   
 $6y = -3x + 1$        $m = \frac{1}{2}$   
 $y = -\frac{1}{2}x + \frac{1}{6}$   
 $m = -\frac{1}{2}$

The lines are neither.

**40.**  $x = 3y + 4$        $y = -3x + 7$   
 $-3y = -x + 4$        $m = -3$   
 $y = \frac{1}{3}x - \frac{4}{3}$   
 $m = \frac{1}{3}$

The lines are perpendicular.

**41.**  $y = 3$        $x = 4$   
 horizontal      vertical  
 The lines are perpendicular.

**42.**  $y = -3$        $y = -7$   
 horizontal      horizontal  
 The lines are parallel.

**43.**  $x = \frac{y-2}{3}$        $3(y-3) + x = 0$   
 $3x = y-2$        $3y-9+x=0$   
 $-y = -3x-2$        $3y = -x+9$   
 $y = 3x+2$        $y = -\frac{1}{3}x+3$   
 $m = 3$        $m = -\frac{1}{3}$

The lines are perpendicular.

**44.**  $2y = 8$        $3(2+x) = 3(y+2)$  /  
 $y = 4$        $6+3x = 3y+6$   
 horizontal       $-3y = -3x$   
 $y = x$   
 $m = 1$

neither

**45.**  $y - y_1 = m(x - x_1)$   
 $y - 4 = 2(x - 2)$   
 $y - 4 = 2x - 4$   
 $-2x + y = 0$   
 $2x - y = 0$

**46.**  $y - y_1 = m(x - x_1)$   
 $y - 5 = -3(x - 3)$   
 $y - 5 = -3x + 9$   
 $3x + y = 14$

**47.**  $y - y_1 = m(x - x_1)$   
 $y - \frac{1}{2} = 2\left(x + \frac{3}{2}\right)$   
 $y - \frac{1}{2} = 2x + 3$   
 $2y - 1 = 4x + 6$   
 $-4x + 2y = 7$   
 $4x - 2y = -7$

**48.**  $y - y_1 = m(x - x_1)$   
 $y + 2 = -6\left(x - \frac{1}{4}\right)$   
 $y + 2 = -6x + \frac{3}{2}$   
 $2y + 4 = -12x + 3$   
 $12x + 2y = -1$

**49.**  $y - y_1 = m(x - x_1)$   
 $y - 1 = \frac{2}{5}(x + 1)$   
 $5(y - 1) = 2(x + 1)$   
 $5y - 5 = 2x + 2$   
 $-2x + 5y = 7$   
 $2x - 5y = -7$

**50.**  $y - y_1 = m(x - x_1)$   
 $y + 3 = -\frac{1}{5}(x + 2)$   
 $5(y + 3) = -(x + 2)$   
 $5y + 15 = -x - 2$   
 $x + 5y = -17$

**51.**  $y - y_1 = m(x - x_1)$   
 $y + 3 = 0(x + 6)$   
 $y + 3 = 0$   
 $y = -3$

**52.**  $y - y_1 = m(x - x_1)$   
 $y - 5 = 0(x - 7)$   
 $y - 5 = 0$   
 $y = 5$

**53.**  $m$  is und  $\Rightarrow$  vertical  
 $x = \text{constant}$   
 $x = -6$

### EXERCISES 2.4

- 54.**  $m$  is und  $\Rightarrow$  vertical  
 $x = \text{constant}$   
 $x = 6$

**55.**  $y - y_1 = m(x - x_1)$   
 $y - 0 = \pi(x - \pi)$   
 $y = \pi x - \pi^2$   
 $-\pi x + y = -\pi^2$   
 $\pi x - y = \pi^2$

**56.**  $y - y_1 = m(x - x_1)$   
 $y - \pi = \pi(x - 0)$   
 $y - \pi = \pi x$   
 $-\pi x + y = \pi$   
 $\pi x - y = -\pi$

- 57.** From the graph,  $m = \frac{2}{3}$  and the line passes through  $(2, 5)$ .

$$\begin{aligned}y - y_1 &= m(x - x_1) \\y - 5 &= \frac{2}{3}(x - 2) \\3(y - 5) &= 3 \cdot \frac{2}{3}(x - 2) \\3y - 15 &= 2(x - 2) \\3y - 15 &= 2x - 4 \\-2x + 3y &= 11 \\2x - 3y &= -11\end{aligned}$$

**59.**  $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{4 - 0}{4 - 0} = \frac{4}{4} = 1$   
 $y - y_1 = m(x - x_1)$   
 $y - 0 = 1(x - 0)$   
 $y = x$

**61.**  $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{-3 - 4}{0 - 3} = \frac{-7}{-3} = \frac{7}{3}$   
 $y - y_1 = m(x - x_1)$   
 $y + 3 = \frac{7}{3}(x - 0)$   
 $y = \frac{7}{3}x - 3$

**63.** From the graph,  $m = -\frac{9}{5}$  and the line passes through  $(-2, 4)$   
 $y - y_1 = m(x - x_1)$   
 $y - 4 = -\frac{9}{5}(x + 2)$   
 $y - 4 = -\frac{9}{5}x - \frac{18}{5}$   
 $y = -\frac{9}{5}x - \frac{18}{5} + 4$   
 $y = -\frac{9}{5}x + \frac{2}{5}$

- 58.** From the graph,  $m = -\frac{2}{3}$  and the line passes through  $(-3, 2)$ .

$$\begin{aligned}y - y_1 &= m(x - x_1) \\y - 2 &= -\frac{2}{3}(x + 3) \\3(y - 2) &= 3 \cdot \left(-\frac{2}{3}\right)(x + 3) \\3y - 6 &= -2(x + 3) \\3y - 6 &= -2x - 6 \\2x + 3y &= 0\end{aligned}$$

**60.**  $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{0 - (-5)}{0 - (-5)} = \frac{5}{5} = 1$   
 $y - y_1 = m(x - x_1)$   
 $y - 0 = 1(x - 0)$   
 $y = x$

**62.**  $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{-8 - 0}{6 - 4} = \frac{-8}{2} = -4$   
 $y - y_1 = m(x - x_1)$   
 $y - 0 = -4(x - 4)$   
 $y = -4x + 16$

**64.** From the graph,  $m = \frac{8}{5}$  and the line passes through  $(2, 3)$ .  
 $y - y_1 = m(x - x_1)$   
 $y - 3 = \frac{8}{5}(x - 2)$   
 $y - 3 = \frac{8}{5}x - \frac{16}{5}$   
 $y = \frac{8}{5}x - \frac{16}{5} + 3$   
 $y = \frac{8}{5}x - \frac{1}{5}$

**EXERCISES 2.4**

**65.**  $y = 4x - 7$      $y - y_1 = m(x - x_1)$   
 $m = 4$                    $y - 0 = 4(x - 0)$   
 Use  $m = 4$ .              y = 4x

**66.**  $x = -3y - 12$      $y - y_1 = m(x - x_1)$   
 $3y = -x - 12$             $y - 0 = -\frac{1}{3}(x - 0)$   
 $y = -\frac{1}{3}x - 4$        y = -\frac{1}{3}x  
 $m = -\frac{1}{3}$   
 Use  $m = -\frac{1}{3}$ .

**67.**  $4x - y = 7$              $y - y_1 = m(x - x_1)$   
 $-y = -4x + 7$             $y - 5 = 4(x - 2)$   
 $y = 4x - 7$                $y - 5 = 4x - 8$   
 $m = 4$                     y = 4x - 3  
 Use  $m = 4$ .

**68.**  $y + 3x = -12$          $y - y_1 = m(x - x_1)$   
 $y = -3x - 12$             $y - 3 = -3(x + 6)$   
 $m = -3$                     $y - 3 = -3x - 18$   
 Use  $m = -3$ .              y = -3x - 15

**69.**  $x = \frac{5}{4}y - 2$      $y - y_1 = m(x - x_1)$   
 $4x = 5y - 8$              $y + 2 = \frac{4}{5}(x - 4)$   
 $-5y = -4x - 8$          $y + 2 = \frac{4}{5}x - \frac{16}{5}$   
 $y = \frac{4}{5}x + \frac{8}{5}$    y =  $\frac{4}{5}x - \frac{26}{5}$   
 $m = \frac{4}{5}$   
 Use  $m = \frac{4}{5}$ .

**70.**  $x = -\frac{3}{4}y + 5$      $y - y_1 = m(x - x_1)$   
 $4x = -3y + 20$          $y + 5 = -\frac{4}{3}(x - 1)$   
 $3y = -4x + 20$          $y + 5 = -\frac{4}{3}x + \frac{4}{3}$   
 $y = -\frac{4}{3}x + \frac{20}{3}$    y =  $-\frac{4}{3}x - \frac{11}{3}$   
 $m = -\frac{4}{3}$   
 Use  $m = -\frac{4}{3}$ .

**71.**  $y = 4x - 7$              $y - y_1 = m(x - x_1)$   
 $m = 4$                      $y - 0 = -\frac{1}{4}(x - 0)$   
 Use  $m = -\frac{1}{4}$ .        y =  $-\frac{1}{4}x$

**72.**  $x = -3y - 12$          $y - y_1 = m(x - x_1)$   
 $3y = -x - 12$             $y - 0 = 3(x - 0)$   
 $y = -\frac{1}{3}x - 4$        y = 3x  
 $m = -\frac{1}{3}$   
 Use  $m = 3$ .

**73.**  $4x - y = 7$              $y - y_1 = m(x - x_1)$   
 $-y = -4x + 7$             $y - 5 = -\frac{1}{4}(x - 2)$   
 $y = 4x - 7$                $y - 5 = -\frac{1}{4}x + \frac{1}{2}$   
 $m = 4$                     y =  $-\frac{1}{4}x + \frac{11}{2}$   
 Use  $m = -\frac{1}{4}$ .

**74.**  $y + 3x = -12$          $y - y_1 = m(x - x_1)$   
 $y = -3x - 12$             $y - 3 = \frac{1}{3}(x + 6)$   
 $m = -3$                     $y - 3 = \frac{1}{3}x + 2$   
 Use  $m = \frac{1}{3}$ .            y =  $\frac{1}{3}x + 5$

### EXERCISES 2.4

**75.**  $x = \frac{5}{4}y - 2$        $y - y_1 = m(x - x_1)$

$$\begin{aligned} 4x &= 5y - 8 \\ -5y &= -4x - 8 \\ y &= \frac{4}{5}x + \frac{8}{5} \\ m &= \frac{4}{5} \end{aligned}$$

Use  $m = -\frac{5}{4}$ .

$$\boxed{y = -\frac{5}{4}x + 3}$$

**76.**  $x = -\frac{3}{4}y + 5$        $y - y_1 = m(x - x_1)$

$$\begin{aligned} 4x &= -3y + 20 \\ 3y &= -4x + 20 \\ y &= -\frac{4}{3}x + \frac{20}{3} \\ m &= -\frac{4}{3} \end{aligned}$$

Use  $m = \frac{3}{4}$ .

$$\boxed{y = \frac{3}{4}x - \frac{23}{4}}$$

- 77.** Since  $y = 3$  is the equation of a horizontal line, any perpendicular line will be vertical. Find the midpoint:

$$x = \frac{2 + (-6)}{2} = -2; y = \frac{4 + 10}{2} = 7$$

The vertical line through  $(-2, 7)$  is  $x = -2$ .

- 79.** Since  $x = 3$  is the equation of a vertical line, any parallel line will be vertical. Find the midpoint:

$$x = \frac{2 + 8}{2} = 5; y = \frac{-4 + 12}{2} = 4$$

The vertical line through  $(5, 4)$  is  $x = 5$ .

- 81.** Let  $x =$  the number of years the truck has been owned and let  $y =$  the value of the truck. Then two points on the line are given:  $(0, 24300)$  and  $(7, 1900)$ .

$$m = \frac{24300 - 1900}{0 - 7} = \frac{22400}{-7} = -3200$$

$$y - y_1 = m(x - x_1)$$

$$y - 24300 = -3200(x - 0)$$

$$y - 24300 = -3200x$$

$$y = -3200x + 24300$$

- 78.** Since  $y = -8$  is the equation of a horizontal line, any parallel line will be horizontal. Find the midpoint:

$$x = \frac{-4 + (-2)}{2} = -3; y = \frac{2 + 8}{2} = 5$$

The horizontal line through  $(-3, 5)$  is  $y = 5$ .

- 80.** Since  $x = 3$  is the equation of a vertical line, any perpendicular line will be horizontal. Find the midpoint:

$$x = \frac{-2 + 4}{2} = 1; y = \frac{2 + (-8)}{2} = -3$$

The horizontal line through  $(1, -3)$ :  $y = -3$ .

- 82.** Let  $x =$  the number of years the laptop has been owned and let  $y =$  the value of the laptop. Then two points on the line are given:  $(0, 2700)$  and  $(4, 300)$ .

$$m = \frac{2700 - 300}{0 - 4} = \frac{2400}{-4} = -600$$

$$y - y_1 = m(x - x_1)$$

$$y - 2700 = -600(x - 0)$$

$$y - 2700 = -600x$$

$$y = -600x + 2700$$

## EXERCISES 2.4

- 83.** Let  $x$  = the number of years the building has been owned and let  $y$  = the value of the building. Then two points on the line are given:  $(0, 475000)$  and  $(10, 950000)$ .

$$m = \frac{950000 - 475000}{10 - 0} = \frac{475000}{10} = 47500$$

$$\begin{aligned}y - y_1 &= m(x - x_1) \\y - 475000 &= 47500(x - 0) \\y - 475000 &= 47500x \\y &= 47500x + 475000\end{aligned}$$

- 85.** Let  $x$  = the number of years the TV has been owned and let  $y$  = the value of the TV. Then two points on the line are given:  $(0, 1900)$  and  $(3, 1190)$ .

$$\begin{aligned}m &= \frac{1900 - 1190}{0 - 3} = \frac{710}{-3} = -\frac{710}{3} \\y - y_1 &= m(x - x_1) \\y - 1900 &= -\frac{710}{3}(x - 0) \\y - 1900 &= -\frac{710}{3}x \\y &= -\frac{710}{3}x + 1900\end{aligned}$$

- 87.** Let  $x$  = the number of years the copier has been owned and let  $y$  = the value of the copier. Then one point on the line is given:  $(0, 1050)$ . Since the copier depreciates by \$120 per year,  $m = -120$ .

$$\begin{aligned}y - y_1 &= m(x - x_1) \\y - 1050 &= -120(x - 0) \\y - 1050 &= -120x \\y &= -120x + 1050\end{aligned}$$

Let  $x = 8$  and find the value of  $y$ :

$$\begin{aligned}y &= -120x + 1050 \\&= -120(8) + 1050 = 90\end{aligned}$$

The salvage value will be \$90.

- 84.** Let  $x$  = the number of years the house has been owned and let  $y$  = the value of the house. Then two points on the line are given:  $(0, 112000)$  and  $(12, 224000)$ .

$$\begin{aligned}m &= \frac{224000 - 112000}{12 - 0} = \frac{112000}{12} = \frac{28000}{3} \\y - y_1 &= m(x - x_1) \\y - 112000 &= \frac{28000}{3}(x - 0) \\y - 112000 &= \frac{28000}{3}x \\y &= \frac{28000}{3}x + 112000\end{aligned}$$

- 86.** Let  $x$  = the number of years the radio has been owned and let  $y$  = the value of the radio. Then two points on the line are given:  $(0, 555)$  and  $(5, 80)$ .

$$\begin{aligned}m &= \frac{555 - 80}{0 - 5} = \frac{475}{-5} = -95 \\y - y_1 &= m(x - x_1) \\y - 555 &= -95(x - 0) \\y - 555 &= -95x \\y &= -95x + 555\end{aligned}$$

Let  $x = 3$  and find the value of  $y$ :

$$\begin{aligned}y &= -95x + 555 \\&= -95(3) + 555 = 270\end{aligned}$$

It will be worth \$270.

- 88.** Let  $x$  = the number of years the jet ski has been owned and let  $y$  = its value. Then two points on the line are given:  $(0, 13800)$  and  $(6, 0)$ .

$$m = \frac{13800 - 0}{0 - 6} = \frac{13800}{-6} = -2300$$

The jet ski depreciates at a rate of \$2300 per year.

## EXERCISES 2.4

- 89.** Let  $x$  = the number of years the table has been owned and let  $y$  = the value of the table. Then one point on the line is given: (2, 450). Since the table appreciates by \$40 per year,  $m = 40$ .

$$y - y_1 = m(x - x_1)$$

$$y - 450 = 40(x - 2)$$

$$y - 450 = 40x - 80$$

$$y = 40x + 370$$

Let  $x = 13$  and find the value of  $y$ :

$$y = 40x + 370$$

$$= 40(13) + 370 = 890$$

The value will be \$890.

- 90.** Let  $x$  = the number of years the clock has been owned and let  $y$  = the value of the clock. Then two points on the line are given: (2, 350) and (5, 530).

$$m = \frac{530 - 350}{5 - 2} = \frac{180}{3} = 60$$

$$y - y_1 = m(x - x_1)$$

$$y - 350 = 60(x - 2)$$

$$y - 350 = 60x - 120$$

$$y = 60x + 230$$

Let  $x = 7$  and find the value of  $y$ :

$$y = 60x + 230$$

$$= 60(7) + 230 = 650$$

It will be worth \$650.

- 91.** Let  $x$  = the number of years the cottage has been owned and let  $y$  = the value of the cottage. Then one point on the line is given: (3, 47700). Since the cottage appreciates by \$3500 per year,  $m = 3500$ .

$$y - y_1 = m(x - x_1)$$

$$y - 47700 = 3500(x - 3)$$

$$y - 47700 = 3500x - 10500$$

$$y = 3500x + 37200$$

Let  $x = 0$  and find the value of  $y$ :

$$y = 3500x + 37200$$

$$= 3500(0) + 37200 = 37200$$

The purchase price was \$37,200.

- 93.** Let  $x$  = the hours of labor and let  $y$  = the labor charge. Then  $m$  = the hourly charge.

$$y = mx \quad y = 46x$$

$$69 = m(1.5) \quad y = 46(5) = 230$$

$$46 = m \quad \text{The charge will be } \$230.$$

- 92.** Let  $x$  = the number of hours of service needed and let  $y$  = the total charge. Then two points on the line are given: (2, 70) and (4, 105)

$$m = \frac{105 - 70}{4 - 2} = \frac{35}{2} = 17.50$$

$$y - y_1 = m(x - x_1)$$

$$y - 70 = 17.50(x - 2)$$

$$y - 70 = 17.50x - 35$$

$$y = 17.50x + 35$$

The hourly charge is \$17.50.

- 94.** Let  $x$  = the number of hundreds of copies and let  $y$  = the total charge. Then  $m$  = the charge per copy and  $b$  = the fixed charge.

$$y = mx + b \quad y = x + 45$$

$$y = 1x + b \quad y = 10 + 45 = 55$$

$$52 = 1(7) + b \quad \text{The charge will be } \$55.$$

$$45 = b$$

## EXERCISES 2.4

- 95.** Let  $x$  = the number of fires and let  $y$  = the population. Then two points on the line are given:  $(300, 57000)$  and  $(325, 59000)$ .

$$m = \frac{59000 - 57000}{325 - 300} = \frac{2000}{25} = 80$$

$$y - y_1 = m(x - x_1)$$

$$y - 57000 = 80(x - 300)$$

$$y - 57000 = 80x - 24000$$

$$y = 80x + 33000$$

Let  $y = 100000$  and find the value of  $x$ :

$$y = 80x + 33000$$

$$100000 = 80x + 33000$$

$$67000 = 80x$$

$837.5 = x \Rightarrow$  There will be about 838 fires when the population is 100,000.

- 97.** Let  $F$  replace  $x$  and  $C$  replace  $y$ . Then two points on the line are given:

$$(32, 0) \text{ and } (212, 100).$$

$$m = \frac{100 - 0}{212 - 32} = \frac{100}{180} = \frac{5}{9}$$

$$C - C_1 = m(F - F_1)$$

$$C - 0 = \frac{5}{9}(F - 32)$$

$$C = \frac{5}{9}(F - 32)$$

- 99.** Let  $y$  = the percent who smoke and let  $x$  = the # of years since 1974. Two points are given:  $(0, 47)$  and  $(20, 29)$ .

$$m = \frac{29 - 47}{20 - 0} = \frac{-18}{20} = -\frac{9}{10}$$

$$y - y_1 = m(x - x_1)$$

$$y - 47 = -\frac{9}{10}(x - 0)$$

$$y = -\frac{9}{10}x + 47$$

Let  $x = 50$ :

$$y = -\frac{9}{10}(50) + 47 = -45 + 47 = 2$$

2% will smoke in 2024.

- 96.** Let  $x$  = the number of feet of gutter and let  $y$  = the total charge. Then  $m$  = the charge per foot. One point on the line is given:  $(250, 435)$

$$y = mx + b$$

$$435 = m(250) + 60$$

$$375 = 250m$$

$$1.5 = m$$

Let  $x = 300$  and find the value of  $y$ :

$$y = 1.5x + 60$$

$$= 1.5(300) + 60 = 510$$

It will cost \$510.

- 98.** Two points on the line are given:

$$(1, 88) \text{ and } (0, 0).$$

$$m = \frac{88 - 0}{1 - 0} = \frac{88}{1} = 88$$

$$y - y_1 = m(x - x_1)$$

$$y - 0 = 88(x - 0)$$

$$y = 88x$$

- 100.** Let  $f$  replace  $x$  and  $h$  replace  $y$ . Then two points on the line are given:

$$(62.5, 200) \text{ and } (40.2, 150).$$

$$m = \frac{150 - 200}{40.2 - 62.5} = \frac{-50}{-22.3} \approx 2.242$$

$$h - h_1 = m(f - f_1)$$

$$h - 200 = 2.242(f - 62.5)$$

$$h - 200 = 2.242f - 140.125$$

$$h = 2.242f + 59.875$$

Let  $f = 50$ :

$$h = 2.242(50) + 59.875 \approx 172$$

He would be about 172 cm tall.

## EXERCISES 2.4

- 101.** Two points on the line are given:

$(0, 37.5)$  and  $(2, 45)$ .

$$m = \frac{45 - 37.5}{2 - 0} = \frac{7.5}{2} = 3.75$$

$$y - y_1 = m(x - x_1)$$

$$y - 37.5 = 3.75(x - 0)$$

$$y = 3.75x + 37.5$$

Let  $x = 10$  and find the value of  $y$ :

$$y = 3.75x + 37.5$$

$$= 3.75(10) + 37.5$$

$$= 37.5 + 37.5 = 75$$

The price will be \$75 in the year 2020.

- 103.** The equation describing the production is  $y = -70x + 1900$ , where  $x$  represents the number of years and  $y$  is the level


of production. Let  $x = 3\frac{1}{2} = \frac{7}{2}$ .

$$y = -70x + 1900$$

$$= -70\left(\frac{7}{2}\right) + 1900 = 1655$$

The production will be 1655 barrels per day.

- 105. a.**



- 102.** Let January be represented by  $x = 0$ , and later months by  $x = 1, 2, 3, \dots$ . Let  $y$  represent the inventory. Then two points on the line are given:  $(0, 375)$  and  $(3, 264)$ .

$$m = \frac{375 - 264}{0 - 3} = \frac{111}{-3} = -37$$

$$y - y_1 = m(x - x_1)$$

$$y - 375 = -37(x - 0)$$

$$y = -37x + 375$$

Let  $x = 2$  and find the value of  $y$ :

$$y = -37x + 375$$

$$= -37(2) + 375 = 301$$

The March inventory will be about 301.

- 104.** Let  $x =$  the number of years the piping has been owned and let  $y =$  the value of the piping. Then two points on the line are given:  $(0, 137000)$  and  $(12, -33000)$ .

$$m = \frac{-33000 - 137000}{12 - 0} = -\frac{42500}{3}$$

$$y - y_1 = m(x - x_1)$$

$$y - 137000 = -\frac{42500}{3}(x - 0)$$

$$y = -\frac{42500}{3}x + 137000$$

- b.** Use  $(50, 20)$  and  $(100, 250)$  for the regression line.

$$m = \frac{250 - 20}{100 - 50} = \frac{230}{50} = \frac{23}{5}$$

$$y - y_1 = m(x - x_1)$$

$$y - 20 = \frac{23}{5}(x - 50)$$

$$y - 20 = \frac{23}{5}x - 230$$


$$y = \frac{23}{5}x - 210$$

$$\text{c. } y = \frac{23}{5}(90) - 210 = 204$$

The rate will be about 204 chirps per minute.

### EXERCISES 2.4

**106. a.**



**b.** Use  $(26, 5)$  and  $(38, 19)$  for the regression line.

$$m = \frac{19 - 5}{38 - 26} = \frac{14}{12} = \frac{7}{6}$$

$$y - y_1 = m(x - x_1)$$

$$y - 5 = \frac{7}{6}(x - 26)$$

$$y - 5 = \frac{7}{6}x - \frac{91}{3}$$

$$y = \frac{7}{6}x - \frac{76}{3}$$

**c.**  $y = \frac{7}{6}(32) - \frac{76}{3} = 12$

The weight will be about 12 pounds.

**107.**  $y = 4.44x - 196.62$

**108.**  $y = 0.96x - 19.22$

**109-112. Answers may vary.**

$$\begin{aligned} \text{113. } m &= \frac{b - 0}{0 - a} = -\frac{b}{a} \\ y - y_1 &= m(x - x_1) \\ y - b &= -\frac{b}{a}(x - 0) \\ y - b &= -\frac{b}{a}x \\ ay - ab &= -bx \\ bx + ay &= ab \\ \frac{bx + ay}{ab} &= \frac{ab}{ab} \\ \frac{x}{a} + \frac{y}{b} &= 1 \end{aligned}$$

**119.**  $Ax + By = C$

$$By = -Ax + C$$

$$y = -\frac{A}{B}x + \frac{C}{B}$$

False.  $m = -\frac{A}{B}$

**121.** Both are horizontal. True.

**123.**  $x = 99$  is vertical, so the parallel line must be vertical too ( $x = -99$ ). False.

**125.**  $\sqrt{5}x + \sqrt{10}y = \sqrt{15}$

$$\frac{\sqrt{5}x + \sqrt{10}y}{\sqrt{5}} = \frac{\sqrt{15}}{\sqrt{5}}$$

$$x + \sqrt{2}y = \sqrt{3}; \text{ True.}$$

$$\begin{array}{ll} \text{114. } & \begin{array}{ll} x\text{-intercept} & y\text{-intercept} \\ bx + ay = ab & bx + ay = ab \\ bx + a(0) = ab & b(0) + ay = ab \\ bx = ab & ay = ab \\ x = a & y = b \\ (a, 0) & (0, b) \end{array} \end{array}$$

**115-118. Answers may vary.**

**120.**  $Ax + By = C$

$$By = -Ax + C$$

$$y = -\frac{A}{B}x + \frac{C}{B}$$

$y$ -intercept:  $\frac{C}{B}$

**122.**  $\frac{\sqrt{11}}{11} \cdot \left(-\sqrt{11}\right) = -\frac{11}{11} = -1$ ; True.

**124.**  $y = 99$  is horizontal, so the perpendicular line must be vertical too ( $x = 99$ ). True.

**126.** False. You can tell by calculating the slopes.

## EXERCISES 2.5

### Exercises 2.5 (page 265)

1.  $x$ -intercept

2.  $y$ -axis

3. axis of symmetry

4.  $y$ -axis

5.  $x$ -axis

6. origin

7. circle, center

8. radius

9.  $x^2 + y^2 = r^2$

10.  $(x - h)^2 + (y - k)^2 = r^2$

11.  $y = x^2 - 4$

$y = x^2 - 4$

12.  $y = x^2 - 9$

$y = x^2 - 9$

$0 = (x + 2)(x - 2)$

$y = 0^2 - 4$

$0 = x^2 - 9$

$y = 0^2 - 9$

$x = -2, x = 2$

$y = -4$

$0 = (x + 3)(x - 3)$

$y = -9$

$x\text{-int: } (-2, 0), (2, 0)$

$y\text{-int: } (0, -4)$

$x = -3, x = 3$

$y\text{-int: } (0, -9)$

$x\text{-int: } (-3, 0), (3, 0)$

13.  $y = 4x^2 - 2x$

$y = 4x^2 - 2x$

14.  $y = 2x - 4x^2$

$y = 2x - 4x^2$

$0 = 2x(2x - 1)$

$y = 4(0)^2 - 2(0)$

$0 = 2x(1 - 2x)$

$y = 2(0) - 4(0)^2$

$x = 0, x = \frac{1}{2}$

$y = 0$

$x = 0, x = \frac{1}{2}$

$y = 0$

$x\text{-int: } (0, 0), (\frac{1}{2}, 0)$

$y\text{-int: } (0, 0)$

$x\text{-int: } (0, 0), (\frac{1}{2}, 0)$

$y\text{-int: } (0, 0)$

15.  $y = x^2 - 4x - 5$

$y = x^2 - 4x - 5$

16.  $y = x^2 - 10x + 21$

$y = x^2 - 10x + 21$

$0 = (x + 1)(x - 5)$

$y = 0^2 - 4(0) - 5$

$0 = (x - 3)(x - 7)$

$y = 0^2 - 10(0) + 21$

$x = -1, x = 5$

$y = -5$

$x = 3, x = 7$

$y = 21$

$x\text{-int: } (-1, 0), (5, 0)$

$y\text{-int: } (0, -5)$

$x\text{-int: } (3, 0), (7, 0)$

$y\text{-int: } (0, 21)$

17.  $y = x^2 + x - 2$

$y = x^2 + x - 2$

18.  $y = x^2 + 2x - 3$

$y = x^2 + 2x - 3$

$0 = (x + 2)(x - 1)$

$y = 0^2 + 0 - 2$

$0 = (x + 3)(x - 1)$

$y = 0^2 + 2(0) - 3$

$x = -2, x = 1$

$y = -2$

$x = -3, x = 1$

$y = -3$

$x\text{-int: } (-2, 0), (1, 0)$

$y\text{-int: } (0, -2)$

$x\text{-int: } (-3, 0), (1, 0)$

$y\text{-int: } (0, -3)$

19.  $y = x^3 - 9x$

$y = x^3 - 9x$

20.  $y = x^3 + x$

$y = x^3 + x$

$0 = x(x^2 - 9)$

$y = 0^3 - 9(0)$

$0 = x(x + 3)(x - 3)$

$y = 0^3 + 0$

$0 = x(x + 3)(x - 3)$

$y = 0$

$x = 0, \{x^2 + 1 \neq 0\}$

$y = 0$

$x = 0, x = -3, x = 3$

$y\text{-int: } (0, 0)$

$x\text{-int: } (0, 0)$

$y\text{-int: } (0, 0)$

$x\text{-int: } (0, 0), (-3, 0), (3, 0)$

21.  $y = x^4 - 1$

$y = x^4 - 1$

22.  $y = x^4 - 25x^2$

$y = x^4 - 25x^2$

$0 = (x^2 + 1)(x^2 - 1)$

$y = 0^4 - 1$

$0 = x^2(x + 5)(x - 5)$

$y = 0$

$0 = (x^2 + 1)(x + 1)(x - 1)$

$y = -1$

$x = 0, x = -5, x = 5$

$y\text{-int: } (0, 0)$

$\{x^2 + 1 \neq 0\}$

$y\text{-int: } (0, -1)$

$x\text{-int: } (0, 0), (-5, 0), (5, 0)$

$x = -1, x = 1$

$y\text{-int: } (0, 0)$


$x\text{-int: } (-1, 0), (1, 0)$

### EXERCISES 2.5

**23.**  $y = x^2$

$x$ -int:  $(0, 0)$


$y$ -int:  $(0, 0)$



**24.**  $y = -x^2$

$x$ -int:  $(0, 0)$


$y$ -int:  $(0, 0)$



**25.**  $y = -x^2 + 2$

$x$ -int:  $(\sqrt{2}, 0), (-\sqrt{2}, 0)$


$y$ -int:  $(0, 2)$



**26.**  $y = x^2 - 1$

$x$ -int:  $(1, 0), (-1, 0)$


$y$ -int:  $(0, -1)$



**27.**  $y = x^2 - 4x$

$x$ -int:  $(0, 0), (4, 0)$


$y$ -int:  $(0, 0)$



**28.**  $y = x^2 + 2x$

$x$ -int:  $(0, 0), (-2, 0)$


$y$ -int:  $(0, 0)$



**29.**  $y = \frac{1}{2}x^2 - 2x$

$x$ -int:  $(0, 0), (4, 0)$


$y$ -int:  $(0, 0)$



**30.**  $y = \frac{1}{2}x^2 + 3$

$x$ -int: none

$y$ -int:  $(0, 3)$



### EXERCISES 2.5

<b>31.</b> $x\text{-axis}$ <hr/> $-y = x^2 + 2$ not equivalent: no symmetry	$y = x^2 + 2$ $y\text{-axis}$ <hr/> $y = (-x)^2 + 2$ $y = x^2 + 2$ equivalent: <span style="border: 1px solid black; padding: 2px;">symmetry</span>	$\text{origin}$ <hr/> $-y = (-x)^2 + 2$ $-y = x^2 + 2$ not equivalent: no symmetry
<b>32.</b> $x\text{-axis}$ <hr/> $-y = 3x + 2$ not equivalent: no symmetry	$y = 3x + 2$ $y\text{-axis}$ <hr/> $y = 3(-x) + 2$ $y = -3x + 2$ not equivalent: no symmetry	$\text{origin}$ <hr/> $-y = 3(-x) + 2$ $-y = -3x + 2$ $y = 3x - 2$ not equivalent: no symmetry
<b>33.</b> $x\text{-axis}$ <hr/> $(-y)^2 + 1 = x$ $y^2 + 1 = x$ equivalent: <span style="border: 1px solid black; padding: 2px;">symmetry</span>	$y^2 + 1 = x$ $y\text{-axis}$ <hr/> $y^2 + 1 = -x$ not equivalent: no symmetry	$\text{origin}$ <hr/> $(-y)^2 + 1 = -x$ $y^2 + 1 = -x$ not equivalent: no symmetry
<b>34.</b> $x\text{-axis}$ <hr/> $(-y)^2 + (-y) = x$ $y^2 - y = x$ not equivalent: no symmetry	$y^2 + y = x$ $y\text{-axis}$ <hr/> $y^2 + y = -x$ not equivalent: no symmetry	$\text{origin}$ <hr/> $(-y)^2 + (-y) = -x$ $y^2 - y = -x$ $-y^2 + y = x$ not equivalent: no symmetry
<b>35.</b> $x\text{-axis}$ <hr/> $(-y)^2 = x^2$ $y^2 = x^2$ equivalent: <span style="border: 1px solid black; padding: 2px;">symmetry</span>	$y^2 = x^2$ $y\text{-axis}$ <hr/> $y^2 = (-x)^2$ $y^2 = x^2$ equivalent: <span style="border: 1px solid black; padding: 2px;">symmetry</span>	$\text{origin}$ <hr/> $(-y)^2 = (-x)^2$ $y^2 = x^2$ equivalent: <span style="border: 1px solid black; padding: 2px;">symmetry</span>
<b>36.</b> $x\text{-axis}$ <hr/> $-y = 3x + 7$ not equivalent: no symmetry	$y = 3x + 7$ $y\text{-axis}$ <hr/> $y = 3(-x) + 7$ $y = -3x + 7$ not equivalent: no symmetry	$\text{origin}$ <hr/> $-y = 3(-x) + 7$ $-y = -3x + 7$ $y = 3x - 7$ not equivalent: no symmetry

### EXERCISES 2.5

**37.**

$x\text{-axis}$	$y = 3x^2 + 7$	$y\text{-axis}$	$\text{origin}$
$-y = 3x^2 + 7$	$y = 3(-x)^2 + 7$	$y = 3x^2 + 7$	$-y = 3(-x)^2 + 7$
not equivalent: no symmetry	equivalent: <span style="border: 1px solid black; padding: 2px;">symmetry</span>	not equivalent: no symmetry	not equivalent: no symmetry

**38.**

$x\text{-axis}$	$x^2 + y^2 = 1$	$y\text{-axis}$	$\text{origin}$
$x^2 + (-y)^2 = 1$	$(-x)^2 + y^2 = 1$	$x^2 + y^2 = 1$	$(-x)^2 + (-y)^2 = 1$
$x^2 + y^2 = 1$	$x^2 + y^2 = 1$	$x^2 + y^2 = 1$	$x^2 + y^2 = 1$

equivalent: symmetry

equivalent: symmetry

equivalent: symmetry

**39.**

$x\text{-axis}$	$y = 3x^3 + 7$	$y\text{-axis}$	$\text{origin}$
$-y = 3x^3 + 7$	$y = 3(-x)^3 + 7$	$y = -3x^3 + 7$	$-y = 3(-x)^3 + 7$
not equivalent: no symmetry			

not equivalent: no symmetry

not equivalent: no symmetry

not equivalent: no symmetry

**40.**

$x\text{-axis}$	$y = 3x^3 + 7x$	$y\text{-axis}$	$\text{origin}$
$-y = 3x^3 + 7x$	$y = 3(-x)^3 + 7x$	$y = -3x^3 + 7x$	$-y = 3(-x)^3 + 7(-x)$
not equivalent: no symmetry	not equivalent: no symmetry	not equivalent: no symmetry	equivalent: <span style="border: 1px solid black; padding: 2px;">symmetry</span>

not equivalent: no symmetry

not equivalent: no symmetry

equivalent: symmetry

**41.**

$x\text{-axis}$	$y^2 = 3x$	$y\text{-axis}$	$\text{origin}$
$(-y)^2 = 3x$	$y^2 = 3(-x)$	$y^2 = -3x$	$(-y)^2 = 3(-x)$
$y^2 = 3x$	$y^2 = -3x$	not equivalent: no symmetry	not equivalent: no symmetry

equivalent: symmetry

not equivalent: no symmetry

not equivalent: no symmetry

**42.**

$x\text{-axis}$	$y = 3x^4 + 7$	$y\text{-axis}$	$\text{origin}$
$-y = 3x^4 + 7$	$y = 3(-x)^4 + 7$	$y = 3x^4 + 7$	$-y = 3(-x)^4 + 7$
not equivalent: no symmetry	equivalent: <span style="border: 1px solid black; padding: 2px;">symmetry</span>	not equivalent: no symmetry	not equivalent: no symmetry

not equivalent: no symmetry

equivalent: symmetry

not equivalent: no symmetry

### EXERCISES 2.5

**43.**

$x\text{-axis}$	$y =  x $	origin
$-y =  x $	$y =  -x $	$-y =  -x $
not equivalent: no symmetry	$y =  -1  x $	$-y =  -1  x $

$$y = |x|$$

$y\text{-axis}$

$$\begin{aligned} y &= |-x| \\ y &= | -1 ||x| \\ y &= |x| \end{aligned}$$

equivalent: symmetry      not equivalent: no symmetry

**44.**

$x\text{-axis}$	$y =  x + 1 $	origin
$-y =  x + 1 $	$y =  -x + 1 $	$-y =  -x + 1 $

$$\begin{aligned} y &= |x + 1| \\ y &= |-x + 1| \\ y &= |-x + 1| \end{aligned}$$

not equivalent: no symmetry      not equivalent: no symmetry      not equivalent: no symmetry

**45.**

$x\text{-axis}$	$ y  = x$	origin
$ -y  = x$	$ y  = -x$	$ -y  = -x$
$ -1  y  = x$	not equivalent: no symmetry	$ -1  y  = -x$
$ y  = x$	$ y  = -x$	$ y  = -x$

equivalent: symmetry      not equivalent: no symmetry

**46.**

$x\text{-axis}$	$ y  =  x $	origin
$ -y  =  x $	$ y  =  -x $	$ -y  =  -x $
$ -1  y  =  x $	$ y  =  -1  x $	$ -1  y  =  -1  x $
$ y  =  x $	$ y  =  x $	$ y  =  x $


equivalent: symmetry      equivalent: symmetry      equivalent: symmetry

**47.**  $y = x^2 + 4x$

$x\text{-int: } (0, 0), (-4, 0)$

$y\text{-int: } (0, 0)$

symmetry: none




**48.**  $y = x^2 - 6x$

$x\text{-int: } (0, 0), (6, 0)$

$y\text{-int: } (0, 0)$

symmetry: none




**49.**  $y = x^3$

$x\text{-int: } (0, 0)$

$y\text{-int: } (0, 0)$

symmetry: origin




## EXERCISES 2.5

50.  $y = x^3 + x$

$x$ -int:  $(0, 0)$

$y$ -int:  $(0, 0)$

symmetry: origin




51.  $y = |x - 2|$

$x$ -int:  $(2, 0)$

$y$ -int:  $(0, 2)$

symmetry: none




52.  $y = |x| - 2$

$x$ -int:  $(-2, 0), (2, 0)$

$y$ -int:  $(0, -2)$

symmetry:  $y$ -axis




53.  $y = -|x| + 3$

$x$ -int:  $(-3, 0), (3, 0)$

$y$ -int:  $(0, 3)$

symmetry:  $y$ -axis




54.  $y = 3|x|$

$x$ -int:  $(0, 0)$

$y$ -int:  $(0, 0)$

symmetry:  $y$ -axis




55.  $y^2 = -x$

$x$ -int:  $(0, 0)$

$y$ -int:  $(0, 0)$

symmetry:  $x$ -axis




56.  $y^2 = 4x$

$x$ -int:  $(0, 0)$

$y$ -int:  $(0, 0)$

symmetry:  $x$ -axis




57.  $y^2 = 9x$

$x$ -int:  $(0, 0)$

$y$ -int:  $(0, 0)$

symmetry:  $x$ -axis




58.  $y^2 = -4x$

$x$ -int:  $(0, 0)$


$y$ -int:  $(0, 0)$

symmetry:  $x$ -axis




### EXERCISES 2.5


**59.**  $y = \sqrt{x} - 1$   
 $x\text{-int: } (1, 0)$   
 $y\text{-int: } (0, -1)$   
 symmetry: none




**60.**  $y = 1 - \sqrt{x}$   
 $x\text{-int: } (1, 0)$   
 $y\text{-int: } (0, 1)$   
 symmetry: none




**61.**  $xy = 4$   
 $x\text{-int: none}$   
 $y\text{-int: none}$   
 symmetry: origin




**62.**  $xy = -9$   
 $x\text{-int: none}$   
 $y\text{-int: none}$   
 symmetry: origin



**63.**  $y = \sqrt[3]{x}$   
 $x\text{-int: } (0, 0)$   
 $y\text{-int: } (0, 0)$   
 symmetry: origin



**64.**  $y = -\sqrt[3]{x}$   
 $x\text{-int: } (0, 0)$   
 $y\text{-int: } (0, 0)$   
 symmetry: origin



**65.**  $x^2 + y^2 = 100$   
 $(x - 0)^2 + (y - 0)^2 = 10^2$   
 C:  $(0, 0); r = 10$

**66.**  $x^2 + y^2 = 81$   
 $(x - 0)^2 + (y - 0)^2 = 9^2$   
 C:  $(0, 0); r = 9$

**67.**  $x^2 + (y - 5)^2 = 49$   
 $(x - 0)^2 + (y - 5)^2 = 7^2$   
 C:  $(0, 5); r = 7$

**68.**  $x^2 + (y + 3)^2 = 8$   
 $(x - 0)^2 + (y - (-3))^2 = (\sqrt{8})^2$   
 $(x - 0)^2 + (y - (-3))^2 = (2\sqrt{2})^2$   
 C:  $(0, -3); r = 2\sqrt{2}$

**69.**  $(x + 6)^2 + y^2 = \frac{1}{4}$   
 $(x - (-6))^2 + (y - 0)^2 = \left(\frac{1}{2}\right)^2$   
 C:  $(-6, 0); r = \frac{1}{2}$

**70.**  $(x - 5)^2 + y^2 = \frac{16}{25}$   
 $(x - 5)^2 + (y - 0)^2 = \left(\frac{4}{5}\right)^2$   
 C:  $(5, 0); r = \frac{4}{5}$

### EXERCISES 2.5

**71.**  $(x - 4)^2 + (y - 1)^2 = 9$   
 $(x - 4)^2 + (y - 1)^2 = 3^2$   
 C:  $(4, 1)$ ;  $r = 3$

**73.**  $(x - \frac{1}{4})^2 + (y + 2)^2 = 45$   
 $(x - \frac{1}{4})^2 + (y - (-2))^2 = (\sqrt{45})^2$   
 $(x - \frac{1}{4})^2 + (y - (-2))^2 = (3\sqrt{5})^2$   
 C:  $(\frac{1}{4}, -2)$ ;  $r = 3\sqrt{5}$

**75.**  $(x - 0)^2 + (y - 0)^2 = 5^2$   
 $x^2 + y^2 = 25$

**77.**  $(x - 0)^2 + (y - (-6))^2 = 6^2$   
 $x^2 + (y + 6)^2 = 36$

**79.**  $(x - 8)^2 + (y - 0)^2 = (\frac{1}{5})^2$   
 $(x - 8)^2 + y^2 = \frac{1}{25}$

**81.**  $(x - (-2))^2 + (y - 12)^2 = 13^2$   
 $(x + 2)^2 + (y - 12)^2 = 169$

**83.**  $x^2 + y^2 = 1^2 \Rightarrow x^2 + y^2 - 1 = 0$

**85.**  $(x - 6)^2 + (y - 8)^2 = 4^2$   
 $x^2 - 12x + 36 + y^2 - 16y + 64 = 16$   
 $x^2 + y^2 - 12x - 16y + 84 = 0$

**87.**  $(x - 3)^2 + (y + 4)^2 = (\sqrt{2})^2$   
 $x^2 - 6x + 9 + y^2 + 8y + 16 = 2$   
 $x^2 + y^2 - 6x + 8y + 23 = 0$

**72.**  $(x + 11)^2 + (y + 7)^2 = 121$   
 $(x - (-11))^2 + (y - (-7))^2 = 11^2$   
 C:  $(-11, -7)$ ;  $r = 11$

**74.**  $(x + \sqrt{5})^2 + (y - 3)^2 = 1$   
 $(x - (-\sqrt{5}))^2 + (y - 3)^2 = (1)^2$   
 C:  $(-\sqrt{5}, 3)$ ;  $r = 1$

**76.**  $(x - 0)^2 + (y - 0)^2 = (\sqrt{3})^2$   
 $x^2 + y^2 = 3$

**78.**  $(x - 0)^2 + (y - 7)^2 = 9^2$   
 $x^2 + (y - 7)^2 = 81$

**80.**  $(x - (-10))^2 + (y - 0)^2 = (\sqrt{11})^2$   
 $(x + 10)^2 + y^2 = 11$

**82.**  $(x - \frac{2}{7})^2 + (y - (-5))^2 = 7^2$   
 $(x - \frac{2}{7})^2 + (y + 5)^2 = 49$

**84.**  $x^2 + y^2 = 4^2 \Rightarrow x^2 + y^2 - 16 = 0$

**86.**  $(x - 5)^2 + (y - 3)^2 = 2^2$   
 $x^2 - 10x + 25 + y^2 - 6y + 9 = 4$   
 $x^2 + y^2 - 10x - 6y + 30 = 0$

**88.**  $(x + 9)^2 + (y - 8)^2 = (2\sqrt{3})^2$   
 $x^2 + 18x + 81 + y^2 - 16y + 64 = 12$   
 $x^2 + y^2 + 18x - 16y + 133 = 0$

### EXERCISES 2.5

**89.** Center:  $x = \frac{3+3}{2} = 3$ ,  $y = \frac{-2+8}{2} = 3$   
 $r = \text{distance from center to endpoint}$

$$\begin{aligned} &= \sqrt{(3-3)^2 + (3-8)^2} = 5 \\ &(x-3)^2 + (y-3)^2 = 5^2 \\ x^2 - 6x + 9 + y^2 - 6y + 9 &= 25 \end{aligned}$$

$$x^2 + y^2 - 6x - 6y - 7 = 0$$

**91.**  $r = \text{distance from center to origin}$

$$\begin{aligned} &= \sqrt{(0-(-3))^2 + (0-4)^2} = 5 \\ &(x+3)^2 + (y-4)^2 = 5^2 \\ x^2 + 6x + 9 + y^2 - 8y + 16 &= 25 \end{aligned}$$

$$x^2 + y^2 + 6x - 8y = 0$$

**93.**  $x^2 + y^2 - 6x + 4y + 4 = 0$

$$x^2 - 6x + y^2 + 4y = -4$$

$$x^2 - 6x + 9 + y^2 + 4y + 4 = -4 + 9 + 4$$

$$(x-3)^2 + (y+2)^2 = 9$$

**95.**  $x^2 + y^2 - 10x - 12y + 57 = 0$

$$x^2 - 10x + y^2 - 12y = -57$$

$$x^2 - 10x + 25 + y^2 - 12y + 36 = -57 + 25 + 36$$

$$(x-5)^2 + (y-6)^2 = 4$$

**96.**  $x^2 + y^2 + 2x + 18y + 57 = 0$

$$x^2 + 2x + y^2 + 18y = -57$$

$$x^2 + 2x + 1 + y^2 + 18y + 81 = -57 + 1 + 81$$

$$(x+1)^2 + (y+9)^2 = 25$$

**97.**  $2x^2 + 2y^2 - 8x - 16y + 22 = 0$

$$x^2 + y^2 - 4x - 8y + 11 = 0$$

$$x^2 - 4x + y^2 - 8y = -11$$

$$x^2 - 4x + 4 + y^2 - 8y + 16 = -11 + 4 + 16$$

$$(x-2)^2 + (y-4)^2 = 9$$

**90.** Center:  $x = \frac{-5+5}{2} = 0$ ,  $y = \frac{-9+9}{2} = 0$

$$r = \text{distance from center to endpoint}$$

$$= \sqrt{(0-5)^2 + (0-9)^2} = \sqrt{106}$$

$$(x-0)^2 + (y-0)^2 = (\sqrt{106})^2$$

$$x^2 + y^2 = 106$$

$$x^2 + y^2 - 106 = 0$$

**92.**  $r = \text{distance from center to origin}$

$$= \sqrt{(0-(-2))^2 + (0-6)^2} = \sqrt{40}$$

$$(x+2)^2 + (y-6)^2 = (\sqrt{40})^2$$

$$x^2 + 4x + 4 + y^2 - 12y + 36 = 40$$

$$x^2 + y^2 + 4x - 12y = 0$$

**94.**  $x^2 + y^2 + 4x - 8y - 5 = 0$

$$x^2 + 4x + y^2 - 8y = 5$$

$$x^2 + 4x + 4 + y^2 - 8y + 16 = 5 + 4 + 16$$


$$(x+2)^2 + (y-4)^2 = 25$$

**EXERCISES 2.5**

98. 
$$\begin{aligned}3x^2 + 3y^2 + 6x - 30y + 3 &= 0 \\x^2 + y^2 + 2x - 10y + 1 &= 0 \\x^2 + 2x + y^2 - 10y &= -1 \\x^2 + 2x + 1 + y^2 - 10y + 25 &= -1 + 1 + 25 \\(x + 1)^2 + (y - 5)^2 &= 25\end{aligned}$$


99.  $x^2 + y^2 - 25 = 0$   
 $x^2 + y^2 = 25$

$C(0, 0), r = 5$




100.  $x^2 + y^2 - 8 = 0$   
 $x^2 + y^2 = 8$


$C(0, 0), r = \sqrt{8} = 2\sqrt{2}$



101.  $(x - 1)^2 + (y + 2)^2 = 4$   
 $C(1, -2), r = 2$



102.  $(x + 1)^2 + (y - 2)^2 = 9$   
 $C(-1, 2), r = 3$



### EXERCISES 2.5


**103.**  $x^2 + y^2 + 2x - 24 = 0$

$$x^2 + 2x + y^2 = 24$$

$$x^2 + 2x + 1 + y^2 = 24 + 1$$

$$(x + 1)^2 + y^2 = 25$$

$$C(-1, 0), r = 5$$




**105.**  $x^2 + y^2 + 4x + 2y - 11 = 0$

$$x^2 + 4x + y^2 + 2y = 11$$

$$x^2 + 4x + 4 + y^2 + 2y + 1 = 11 + 4 + 1$$

$$(x + 2)^2 + (y + 1)^2 = 16$$

$$C(-2, -1), r = 4$$




**104.**  $x^2 + y^2 - 4y = 12$

$$x^2 + y^2 - 4y + 4 = 12 + 4$$

$$x^2 + (y - 2)^2 = 16$$

$$C(0, 2), r = 4$$




**106.**  $x^2 + y^2 - 6x + 2y + 1 = 0$

$$x^2 - 6x + y^2 + 2y = -1$$

$$x^2 - 6x + 9 + y^2 + 2y + 1 = -1 + 9 + 1$$

$$(x - 3)^2 + (y + 1)^2 = 9$$

$$C(3, -1), r = 3$$



### EXERCISES 2.5


**107.**  $9x^2 + 9y^2 - 12y = 5$

$$x^2 + y^2 - \frac{4}{3}y = \frac{5}{9}$$

$$x^2 + y^2 - \frac{4}{3}y + \frac{4}{9} = \frac{5}{9} + \frac{4}{9}$$

$$x^2 + \left(y - \frac{2}{3}\right)^2 = 1$$

$$C\left(0, \frac{2}{3}\right), r = 1$$




**109.**  $4x^2 + 4y^2 - 4x + 8y + 1 = 0$

$$x^2 + y^2 - x + 2y = -\frac{1}{4}$$

$$x^2 - x + \frac{1}{4} + y^2 + 2y + 1 = -\frac{1}{4} + \frac{1}{4} + 1$$

$$\left(x - \frac{1}{2}\right)^2 + (y + 1)^2 = 1$$

$$C\left(\frac{1}{2}, -1\right), r = 1$$




**108.**  $4x^2 + 4y^2 + 4y = 15$

$$x^2 + y^2 + y = \frac{15}{4}$$

$$x^2 + y^2 + y + \frac{1}{4} = \frac{15}{4} + \frac{1}{4}$$

$$x^2 + \left(y + \frac{1}{2}\right)^2 = 4$$

$$C\left(0, -\frac{1}{2}\right), r = 2$$




**110.**  $9x^2 + 9y^2 - 6x + 18y + 1 = 0$

$$x^2 + y^2 - \frac{2}{3}x + 2y = -\frac{1}{9}$$

$$x^2 - \frac{2}{3}x + \frac{1}{9} + y^2 + 2y + 1 = -\frac{1}{9} + \frac{1}{9} + 1$$

$$\left(x - \frac{1}{3}\right)^2 + (y + 1)^2 = 1$$


$$C\left(\frac{1}{3}, -1\right), r = 1$$



## EXERCISES 2.5


**111.**  $y = 2x^2 - x + 1$

Vertex:  $(0.25, 0.88)$




**114.**  $y = 2x^2 - 3x + 2$

Vertex:  $(0.75, 0.88)$




**112.**  $y = x^2 + 5x - 6$

Vertex:  $(-2.50, -12.25)$



**113.**  $y = 7 + x - x^2$


Vertex:  $(0.50, 7.25)$



**115.** Graph  $y = x^2 - 7$ .

Find the  $x$ -intercepts.


$$x = -2.65, x = 2.65$$



**116.** Graph  $y = x^2 - 3x + 2$ .

Find the  $x$ -intercepts.


$$x = 1.00, x = 2.00$$



**117.** Graph  $y = x^3 - 3$ .

Find the  $x$ -intercepts.

$$x = 1.44$$




**118.** Graph  $y = 3x^3 - x^2 - x$ .

Find the  $x$ -intercepts.

$$x = -0.43, x = 0,$$

$$x = 0.77$$



**119.** Let  $y = 0$ :

$$y = 64t - 16t^2$$

$$0 = 16t(4 - t)$$

$$t = 0 \text{ or } t = 4$$

It strikes the ground after 4 seconds.


**120.** From #119, the flight lasts 4 seconds. Thus, half the flight is 2 seconds. Let  $t = 2$ :

$$y = 64t - 16t^2$$

$$y = 64(2) - 16(2)^2 = 128 - 64 = 64; \text{ The highest point is 64 feet above ground.}$$

## EXERCISES 2.5

**121.**  $D = 0.08V^2 + 0.9V$ ;



**122.** Refer to the graph for #121.

The  $y$ -coordinate for  $x = 30$  is  $y = 99$ .

The  $y$ -coordinate for  $x = 60$  is  $y = 342$ .

$$342 - 99 = 243$$

At 60 mph, 243 more feet is required to stop than at 30 mph.

**123.**  $r = \frac{12}{2} = 6$

$$(x - 0)^2 + (y - 0)^2 = 6^2$$

$$x^2 + y^2 = 36$$

**125.**  $r = \frac{60}{2} = 30$

$$(x - 0)^2 + (y - 35)^2 = 30^2$$

$$x^2 + (y - 35)^2 = 900$$

**127.**  $r = \sqrt{(10 - 7)^2 + (0 - 4)^2} = 5$

$$(x - 7)^2 + (y - 4)^2 = 5^2$$

$$x^2 - 14x + 49 + y^2 - 8y + 16 = 25$$

$$x^2 + y^2 - 14x - 8y + 40 = 0$$

**128.**

First tire

$$C(12, 12), r = 12$$

$$(x - 12)^2 + (y - 12)^2 = 12^2$$

$$x^2 - 24x + 144 + y^2 - 24y + 144 = 144$$

$$x^2 + y^2 - 24x - 24y + 144 = 0$$

**126.**  $r = \frac{30}{2} = 15$

$$(x - 5)^2 + (y - 10)^2 = 15^2$$

$$(x - 5)^2 + (y - 10)^2 = 225$$

Second tire

$$C(36, 12), r = 12$$

$$(x - 36)^2 + (y - 12)^2 = 12^2$$

$$x^2 - 72x + 1296 + y^2 - 24y + 144 = 144$$

$$x^2 + y^2 - 72x - 24y + 1296 = 0$$

**129-132. Answers may vary.**

**133.**  $x^2 - 4x + y^2 - 6y + 13 = 0$

$$x^2 - 4x + 4 + y^2 - 6y + 9 = -13 + 4 + 9$$

$$(x + 2)^2 + (y - 3)^2 = 0 \Rightarrow \text{a single point}$$

**134.**  $x^2 - 12x + y^2 + 4y + 43 = 0$

$$x^2 - 12x + 36 + y^2 + 4y + 4 = -43 + 36 + 4$$

$$(x - 6)^2 + (y + 2)^2 = -3 \Rightarrow \text{nonexistent}$$

**135.** False. The graphs are symmetric with respect to the  $y$ -axis.

**136.** False. The graphs are symmetric with respect to the origin.

## EXERCISES 2.5

**137.** True.

**138.** False. The line  $y = x$  has symmetry with respect to the origin, but not with respect to either the  $x$ - or  $y$ -axis.

**139.** True.

**140.** True.

**141.** False. The graph is the single point  $(4, -\frac{1}{7})$ .

**142.** True.

## Exercises 2.6 (page 276)

**1.** quotient

**2.** ratios

**3.** means

**4.** extremes

**5.** extremes, means

**6.**  $y = kx$

**7.** inverse

**8.** constant

**9.** joint

**10.**  $x^2, z$

$$\frac{4}{x} = \frac{2}{7}$$

$$\frac{5}{2} = \frac{x}{6}$$

$$4 \cdot 7 = 2 \cdot x$$

$$5 \cdot 6 = x \cdot 2$$

$$28 = 2x$$

$$30 = 2x$$

$$14 = x$$

$$15 = x$$

$$\begin{aligned} \text{13. } \frac{x}{2} &= \frac{3}{x+1} \\ x(x+1) &= 3 \cdot 2 \\ x^2 + x &= 6 \\ x^2 + x - 6 &= 0 \\ (x+3)(x-2) &= 0 \\ x = -3 \text{ or } x &= 2 \end{aligned}$$

$$\begin{aligned} \text{14. } \frac{x+5}{6} &= \frac{7}{8-x} \\ (x+5)(8-x) &= 7 \cdot 6 \\ -x^2 + 3x + 40 &= 42 \\ 0 &= x^2 - 3x + 2 \\ 0 &= (x-2)(x-1) \\ x = 1 \text{ or } x &= 2 \end{aligned}$$

**15.** Let  $x$  = the number of women.

$$\begin{aligned} \frac{3}{5} &= \frac{x}{30} \\ 3 \cdot 30 &= 5 \cdot x \\ 90 &= 5x \\ 18 = x &\Rightarrow \text{There are 18 women.} \end{aligned}$$

**16.** Let  $x$  = the number of bags of lime.

$$\begin{aligned} \frac{3}{7} &= \frac{x}{21} \\ 3 \cdot 21 &= x \cdot 7 \\ 63 &= 7x \\ 9 = x &\Rightarrow 9 \text{ bags of lime should be used.} \end{aligned}$$

**17.**  $y = kx$

$$\begin{aligned} 15 &= k(30) \\ \frac{1}{2} &= k \end{aligned}$$

**18.**  $z = kt$

$$\begin{aligned} 21 &= k(7) \\ 3 &= k \end{aligned}$$

$$\begin{aligned} \text{19. } I &= \frac{k}{R} \\ 50 &= \frac{k}{20} \\ 1000 = k & \end{aligned}$$

### EXERCISES 2.6

**20.**  $R = \frac{k}{I^2}$   
 $100 = \frac{k}{25^2}$   
 $100 = \frac{k}{625}$   
 $62500 = k$

**23.**  $y = kx$      $y = \frac{15}{4}x$   
 $15 = k(4)$      $y = \frac{15}{4} \cdot \frac{7}{5}$   
 $\frac{15}{4} = k$      $y = \frac{21}{4}$

**25.**  $w = \frac{k}{z}$      $w = \frac{30}{z}$   
 $10 = \frac{k}{3}$      $w = \frac{30}{5}$   
 $30 = k$      $w = 6$

**27.**  $P = krs$      $P = -\frac{2}{5}rs$   
 $16 = k(5)(-8)$      $P = -\frac{2}{5}(2)(10)$   
 $16 = -40k$      $P = -8$   
 $-\frac{16}{40} = k$   
 $-\frac{2}{5} = k$

**29.** direct

**30.** neither

**21.**  $E = kIR$

$125 = k(5)(25)$   
 $125 = 125k$   
 $1 = k$

**22.**  $z = k(x + y)$

$28 = k(2 + 5)$   
 $28 = 7k$   
 $4 = k$

**24.**  $w = kz$      $w = -3z$   
 $-6 = k(2)$      $w = -3(-3)$   
 $-3 = k$      $w = 9$

**26.**  $y = \frac{k}{x}$      $y = \frac{200}{x}$   
 $100 = \frac{k}{2}$      $y = \frac{200}{50}$   
 $200 = k$      $y = 4$

**28.**  $m = kn^2\sqrt{q}$      $m = 3n^2\sqrt{q}$   
 $24 = k(2)^2\sqrt{4}$      $m = 3(5)^2\sqrt{9}$   
 $24 = k(4)(2)$      $m = 3(25)(3)$   
 $24 = 8k$      $m = 225$   
 $3 = k$

**32.** inverse

**33.** Let  $x$  = the amount of caffeine.

$$\begin{array}{lll} \frac{55}{12} = \frac{x}{44} & \frac{47}{12} = \frac{x}{44} & \frac{37}{12} = \frac{x}{44} \\ 55 \cdot 44 = 12 \cdot x & 47 \cdot 44 = 12 \cdot x & 37 \cdot 44 = 12 \cdot x \\ 2420 = 12x & 2068 = 12x & 1628 = 12x \\ 202 \text{ mg} \approx x & 172 \text{ mg} \approx x & 136 \text{ mg} \approx x \end{array}$$

**34.** Let  $x$  = the number of phones.

$$\begin{array}{l} \frac{221}{250} = \frac{x}{280000} \\ 221 \cdot 280000 = 250 \cdot x \\ 61880000 = 250x \\ 247,520 = x \\ \text{247,520 have cellular phones.} \end{array}$$

**35.** Let  $x$  = the amount of adhesive needed.

$$\begin{array}{l} \frac{\frac{1}{2}}{140} = \frac{x}{500} \\ \frac{1}{2} \cdot 500 = 140 \cdot x \\ 250 = 140x \\ 1.79 \approx x \\ \text{About 2 gallons of adhesive will be needed.} \end{array}$$

## EXERCISES 2.6

- 36.** Let  $x =$  the dosage.

$$\frac{0.006}{1} = \frac{x}{30}$$

$$0.006 \cdot 30 = 1 \cdot x$$

$$0.18 = x$$

The dosage should be 0.18 g, or 180 mg.

$$\begin{aligned} \text{37. } V &= \frac{kT}{P} & V &= \frac{\frac{80}{33}T}{P} \\ 20 &= \frac{k(330)}{40} & V &= \frac{\frac{80}{33}(300)}{50} \\ 800 &= 330k & V &= \frac{8000}{50} \\ \frac{800}{330} &= k & V &= \frac{160}{11} \\ \frac{80}{33} &= k & V &= 14\frac{6}{11} \text{ ft}^3 \end{aligned}$$

$$\begin{aligned} \text{38. } f &= kd & f &= 25d \\ 5 &= k(0.2) & f &= 25(0.35) \\ 25 &= k & f &= 8.75 \text{ Newtons} \end{aligned}$$

$$\begin{aligned} \text{39. } d &= kt^2 & d &= 16t^2 \\ 16 &= k(1)^2 & 144 &= 16t^2 \\ 16 &= k & 9 &= t^2 \\ & & 3 &= t \Rightarrow 3 \text{ seconds} \end{aligned}$$

$$\begin{aligned} \text{40. } P &= \frac{kV^2}{R} & P &= \frac{V^2}{R} \\ 20 &= \frac{k(20)^2}{20} & 40 &= \frac{V^2}{10} \\ 400 &= 400k & 400 &= V^2 \\ 1 &= k & 20 &= V \Rightarrow 20 \text{ volts} \end{aligned}$$

$$\begin{aligned} \text{41. } t &= kt^2 & t &= l^2 \\ 1 &= k(1)^2 & 2 &= l^2 \\ 1 &= k & \sqrt{2} &= l \Rightarrow \sqrt{2} \text{ meters} \end{aligned}$$

$$\begin{aligned} \text{42. } f &= k\sqrt{T} & f &= \frac{144}{\sqrt{2}}\sqrt{T} \\ 144 &= k\sqrt{2} & f &= \frac{144}{\sqrt{2}}\sqrt{18} \\ \frac{144}{\sqrt{2}} &= k & f &= 144\sqrt{9} \\ & & f &= 144(3) = 432 \text{ hertz} \end{aligned}$$

$$\begin{aligned} \text{43. } I &= \frac{k}{d^2} & I &= \frac{6000}{d^2} \\ 60 &= \frac{k}{10^2} & I &= \frac{6000}{20^2} \\ 60 &= \frac{k}{100} & I &= \frac{6000}{400} \\ 6000 &= k & I &= 15 \Rightarrow 15 \text{ lumens} \end{aligned}$$

$$\begin{aligned} \text{44. } I &= \frac{k}{d^2} & I &= \frac{22500}{d^2} \\ 100 &= \frac{k}{15^2} & I &= \frac{22500}{25^2} \\ 100 &= \frac{k}{225} & I &= \frac{22500}{625} \\ 22500 &= k & I &= 36 \Rightarrow 36 \text{ lumens} \end{aligned}$$

$$\begin{aligned} \text{45. } E &= kmv^2 = k(2m)(3v)^2 \\ &= k(2m)(9v^2) \\ &= 18 \cdot kmv^2 \end{aligned}$$

The energy is multiplied by 18.

$$\begin{aligned} \text{46. } P &= kRC^2 & P &= RC^2 \\ 10 &= k(10)(1)^2 & P &= 5(3)^2 \\ 10 &= 10k & P &= 5(9) \\ 1 &= k & P &= 45 \text{ watts} \end{aligned}$$

### EXERCISES 2.6


47. 
$$\begin{aligned} G &= \frac{km_1m_2}{d^2} = \frac{k(3m_1)(3m_2)}{(2d)^2} \\ &= \frac{k \cdot 9m_1m_2}{4d^2} \\ &= \frac{9}{4} \cdot \frac{km_1m_2}{d^2} \end{aligned}$$

The force is multiplied by  $\frac{9}{4}$ .

48. 
$$\begin{aligned} G &= \frac{km_1m_2}{d^2} = \frac{k(2m_1)(3m_2)}{\left(\frac{d}{2}\right)^2} \\ &= \frac{k \cdot 6m_1m_2}{\frac{d^2}{4}} \\ &= 24 \cdot \frac{km_1m_2}{d^2} \end{aligned}$$

The force is multiplied by 24.


49. Consider this figure:



$h = \sqrt{3}$  can be computed using the Pythagorean Theorem.

$$\begin{aligned} A &= \frac{1}{2}bh = \frac{1}{2}(2)\sqrt{3} = \sqrt{3} \\ A &= ks^2 \\ \sqrt{3} &= k(2)^2 \\ \sqrt{3} &= 4k \\ \frac{\sqrt{3}}{4} &= k \end{aligned}$$

50. Consider this figure:



The diagonal is obtained by repeatedly using the Pythagorean Theorem.

$$\begin{aligned} d &= ks \\ \sqrt{3} &= k(1) \\ \sqrt{3} &= k \end{aligned}$$

51-58. Answers may vary.

59. d

60. c

61. b

62. a

63. c

64. d

65. a

66. b

### Chapter 2 Review (page 279)

1.  $D = \{3, 4, 5, 6\}; R = \{4, 5, 6, 7\}$

Each element of the domain is paired with only one element of the range. Function.

2.  $D = \{2, 3, -4\}; R = \{4, 5, 6, 3\}$

2 is both paired with 4 and 5. Not a function.

## CHAPTER 2 REVIEW

- 3.**  $y = 3$   
Each value of  $x$  is paired with only one value of  $y$ . **function**
- 4.**  $y + 5x^2 = 2$   
 $y = -5x^2 + 2$   
Each value of  $x$  is paired with only one value of  $y$ . **function**
- 5.**  $y^2 - x = 5$   
 $y^2 = x + 5$   
 $y = \pm\sqrt{x+5}$   
Each value of  $x$  is paired with more than one value of  $y$ . **not a function**
- 6.**  $y = |x| + x$   
Each value of  $x$  is paired with only one value of  $y$ . **function**
- 7.**  $f(x) = y = 3x^2 - 5$   
domain =  $(-\infty, \infty)$
- 8.**  $f(x) = y = \frac{3x}{x-5}$   
domain =  $(-\infty, 5) \cup (5, \infty)$
- 9.**  $f(x) = \frac{3x}{4x^2 - 16} = \frac{3x}{4(x+2)(x-2)}$   
 $x \neq -2, x \neq 2$   
domain =  $(-\infty, -2) \cup (-2, 2) \cup (2, \infty)$
- 10.**  $f(x) = y = \sqrt{x-1}$   
domain =  $[1, \infty)$
- 11.**  $f(x) = y = \sqrt{5-x}$   
domain =  $(-\infty, 5]$
- 12.**  $f(x) = y = \sqrt{x^2 + 1}$   
 $x^2 + 1 \geq 0 \Rightarrow$  domain =  $(-\infty, \infty)$
- 13.**  $f(x) = 5x - 2$   
 $f(2) = 5(2) - 2 = 8$   
 $f(-3) = 5(-3) - 2 = -17$   
 $f(k) = 5k - 2$
- 14.**  $f(x) = \frac{6}{x-5}$   
 $f(2) = \frac{6}{2-5} = \frac{6}{-3} = -2$   
 $f(-3) = \frac{6}{-3-5} = \frac{6}{-8} = -\frac{3}{4}$   
 $f(k) = \frac{6}{k-5}$
- 15.**  $f(x) = |x - 2|$   
 $f(2) = |2 - 2| = |0| = 0$   
 $f(-3) = |-3 - 2| = |-5| = 5$   
 $f(k) = |k - 2|$
- 16.**  $f(x) = \frac{x^2 - 3}{x^2 + 3}$   
 $f(2) = \frac{2^2 - 3}{2^2 + 3} = \frac{1}{7}$   
 $f(-3) = \frac{(-3)^2 - 3}{(-3)^2 + 3} = \frac{6}{12} = \frac{1}{2}$   
 $f(k) = \frac{k^2 - 3}{k^2 + 3}$
- 17.** 
$$\begin{aligned} \frac{f(x+h) - f(x)}{h} &= \frac{[5(x+h) - 6] - [5x - 6]}{h} = \frac{[5x + 5h - 6] - [5x - 6]}{h} \\ &= \frac{5x + 5h - 6 - 5x + 6}{h} = \frac{5h}{h} = 5 \end{aligned}$$

## CHAPTER 2 REVIEW

18. 
$$\begin{aligned} \frac{f(x+h) - f(x)}{h} &= \frac{[2(x+h)^2 - 7(x+h) + 3] - [2x^2 - 7x + 3]}{h} \\ &= \frac{[2x^2 + 4xh + 2h^2 - 7x - 7h + 3] - [2x^2 - 7x + 3]}{h} \\ &= \frac{2x^2 + 4xh + 2h^2 - 7x - 7h + 3 - 2x^2 + 7x - 3}{h} \\ &= \frac{4xh + 2h^2 - 7h}{h} = \frac{h(4x + 2h - 7)}{h} = 4x + 2h - 7 \end{aligned}$$

19.  $f(x) = -0.6x + 132$   
 $f(45) = -0.6(45) + 132 = 105$

20. a.  $I(h) = 3.5h - 50$   
 b.  $I(200) = 3.5(200) - 50 = \$650$

21.  $A(2, 0)$

22.  $B(-2, 1)$

23.  $C(0, -1)$


24.  $D(3, -1)$

25.  $A(-3, 5)$ : QII

26.  $B(5, -3)$ : QIV


27.  $C(0, -7)$ : negative  $y$ -axis

28.  $D\left(-\frac{1}{2}, 0\right)$ : negative  $x$ -axis




29.  $2x - y = 6$   
 $-y = -2x + 6$   
 $y = 2x - 6$

$x$	$y$
0	-6
2	-2




30.  $2x + 5y = -10$   
 $5y = -2x - 10$   
 $y = -\frac{2}{5}x - 2$

$x$	$y$
0	-2
-5	0




## CHAPTER 2 REVIEW


**31.**  $3x - 5y = 15$        $3x - 5y = 15$   
 $3x - 5(0) = 15$        $3(0) - 5y = 15$   
 $3x = 15$        $-5y = 15$   
 $x = 5$        $y = -3$   
 $(5, 0)$        $(0, -3)$




**33.**  $x + y = -7$        $x + y = -7$   
 $x + 0 = -7$        $0 + y = -7$   
 $x = -7$        $y = -7$   
 $(-7, 0)$        $(0, -7)$




**35.**  $y = 4 \Rightarrow$  horizontal




**32.**  $x + y = 7$        $x + y = 7$   
 $x + 0 = 7$        $0 + y = 7$   
 $x = 7$        $y = 7$   
 $(7, 0)$        $(0, 7)$



**34.**  $x - 5y = 5$        $x - 5y = 5$   
 $x - 5(0) = 5$        $0 - 5y = 5$   
 $x = 5$        $-5y = 5$   
 $(5, 0)$        $y = -1$   
 $(0, -1)$



**36.**  $x = -2 \Rightarrow$  vertical



**37.** Let  $x = 3$ :  $y = -2200x + 18,750 = -2200(3) + 18,750 = -6600 + 18,750 = \$12,150$

**38.** Let  $x = 5$ :  $y = 16,500x + 250,000 = 16,500(5) + 250,000 = 82,500 + 250,000 = \$332,500$

## CHAPTER 2 REVIEW

- 39.**  $d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$
- $$\begin{aligned} &= \sqrt{(-3 - 3)^2 + (7 - (-1))^2} \\ &= \sqrt{(-6)^2 + (8)^2} \\ &= \sqrt{36 + 64} = \sqrt{100} = 10 \end{aligned}$$
- 40.**  $d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$
- $$\begin{aligned} &= \sqrt{(-12 - (-8))^2 + (10 - 6)^2} \\ &= \sqrt{(-4)^2 + 4^2} \\ &= \sqrt{16 + 16} = \sqrt{32} = 4\sqrt{2} \end{aligned}$$
- 41.**  $d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$
- $$\begin{aligned} &= \sqrt{(\sqrt{3} - \sqrt{3})^2 + (9 - 7)^2} \\ &= \sqrt{0^2 + (2)^2} \\ &= \sqrt{0 + 4} = \sqrt{4} = 2 \end{aligned}$$
- 42.**  $d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$
- $$\begin{aligned} &= \sqrt{(a - (-a))^2 + (-a - a)^2} \\ &= \sqrt{(2a)^2 + (-2a)^2} \\ &= \sqrt{4a^2 + 4a^2} = \sqrt{8a^2} = 2\sqrt{2}|a| \end{aligned}$$
- 43.**  $M\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}\right) = M\left(\frac{-3 + 3}{2}, \frac{7 + (-1)}{2}\right) = M\left(\frac{0}{2}, \frac{6}{2}\right) = M(0, 3)$
- 44.**  $M\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}\right) = M\left(\frac{0 + (-12)}{2}, \frac{5 + 10}{2}\right) = M\left(\frac{-12}{2}, \frac{15}{2}\right) = M\left(-6, \frac{15}{2}\right)$
- 45.**  $M\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}\right) = M\left(\frac{\sqrt{3} + \sqrt{3}}{2}, \frac{9 + 7}{2}\right) = M\left(\frac{2\sqrt{3}}{2}, \frac{16}{2}\right) = M(\sqrt{3}, 8)$
- 46.**  $M\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}\right) = M\left(\frac{a + (-a)}{2}, \frac{-a + a}{2}\right) = M\left(\frac{0}{2}, \frac{0}{2}\right) = M(0, 0)$
- 47.**  $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{7 - (-5)}{1 - 3} = \frac{12}{-2} = -6$
- 48.**  $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{-7 - 7}{-5 - 2} = \frac{-14}{-7} = 2$
- 49.**  $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{\frac{1}{2} - (-8)}{5 - 5} = \frac{\frac{17}{2}}{0}$ : und.
- 50.**  $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{-8 - (-8)}{-1 - \frac{2}{3}} = \frac{0}{-\frac{5}{3}} = 0$
- 51.**  $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{b - a}{a - b} = -1$
- 52.**  $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{(b - a) - b}{b - (a + b)} = \frac{-a}{-a} = 1$
- 53.**  $y = 3x + 6 \quad m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{9 - 6}{1 - 0}$
- |     |     |
|-----|-----|
| $x$ | $y$ |
| 0   | 6   |
| 1   | 9   |
- $$= \frac{3}{1} = 3$$
- 54.**  $y = -\frac{1}{5}x - 6 \quad m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{-7 - (-6)}{5 - 0}$
- |     |     |
|-----|-----|
| $x$ | $y$ |
| 0   | -6  |
| 5   | -7  |
- $$= \frac{-1}{5} = -\frac{1}{5}$$
- 55.** The slope is zero.
- 56.** The slope is undefined.
- 57.** The slope is negative.
- 58.** The slope is positive.

## CHAPTER 2 REVIEW

**59.**  $m_1 m_2 = 5 \left(-\frac{1}{5}\right) = -1$   
perpendicular

**60.**  $m_1 \neq m_2$ ;  $m_1 m_2 = \frac{2}{7} \cdot \frac{7}{2} = 1 \neq -1$   
neither

**61.**  $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{10 - 5}{6 - (-2)} = \frac{5}{8}$   
 $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{y - 2}{10 - 2} = \frac{5}{8}$   
 $8(y - 2) = 5(8)$   
 $8y - 16 = 40$   
 $8y = 56$   
 $y = 7$

**62.**  $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{10 - 5}{6 - (-2)} = \frac{5}{8}$   
 $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{-3 - 5}{x - (-2)} = \frac{-8}{5}$   
 $5(-8) = -8(x + 2)$   
 $-40 = -8x - 16$   
 $8x = 24$   
 $x = 3$

**63.**  $m = \frac{\Delta y}{\Delta x} = \frac{3000}{15} = 200$  ft per minute

**64.**  $m = \frac{\Delta y}{\Delta x} = \frac{147,500 - 50,000}{3 - 1} = \frac{97,500}{2} = \$48,750$  per year

**65.**  $y = mx + b$   
 $y = \frac{2}{3}x + 3$

**66.**  $y = mx + b$   
 $y = -\frac{3}{2}x - 5$

**67.**  $3x - 2y = 10$   
 $-2y = -3x + 10$   
 $y = \frac{3}{2}x - 5$   
 $m = \frac{3}{2}, (0, -5)$

**68.**  $2x + 4y = -8$   
 $4y = -2x - 8$   
 $y = -\frac{1}{2}x - 2$   
 $m = -\frac{1}{2}, (0, -2)$


**69.**  $-2y = -3x + 10$   
 $y = \frac{3}{2}x - 5$   
 $m = \frac{3}{2}, (0, -5)$

**70.**  $2x = -4y - 8$   
 $4y = -2x - 8$   
 $y = -\frac{1}{2}x - 2$   
 $m = -\frac{1}{2}, (0, -2)$


**71.**  $5x + 2y = 7$   
 $2y = -5x + 7$   
 $y = -\frac{5}{2}x + \frac{7}{2}$   
 $m = -\frac{5}{2}, (0, \frac{7}{2})$

**72.**  $3x - 4y = 14$   
 $-4y = -3x + 14$   
 $y = \frac{3}{4}x - \frac{7}{2}$   
 $m = \frac{3}{4}, (0, -\frac{7}{2})$

**73.**  $y = \frac{3}{5}x - 2$   
 $m = \frac{3}{5}, b = -2$



**74.**  $y = -\frac{4}{3}x + 3$   
 $m = -\frac{4}{3}, b = 3$



## CHAPTER 2 REVIEW

**75.**  $y = 3x + 8$      $2y = 6x - 19$   
 $m = 3$                  $y = 3x - \frac{19}{2}$   
 $m = 3$

The lines are parallel.

**76.**  $2x + 3y = 6$                  $3x - 2y = 15$   
 $3y = -2x + 6$                  $-2y = -3x + 15$   
 $y = -\frac{2}{3}x + 2$                  $y = \frac{3}{2}x - \frac{15}{2}$   
 $m = -\frac{2}{3}$                  $m = \frac{3}{2}$

The lines are perpendicular.

**77.**  $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{7 - 0}{-5 - 0} = -\frac{7}{5}$   
 $y - y_1 = m(x - x_1)$   
 $y - 0 = -\frac{7}{5}(x - 0)$   
 $y = -\frac{7}{5}x$   
 $5y = 5\left(-\frac{7}{5}x\right)$   
 $5y = -7x$   
 $7x + 5y = 0$

**78.**  $y - y_1 = m(x - x_1)$   
 $y - 1 = -4(x + 2)$   
 $y - 1 = -4x - 8$   
 $4x + y = -7$

**79.**  $y - y_1 = m(x - x_1)$   
 $y + 1 = -\frac{1}{5}(x - 2)$   
 $5(y + 1) = 5 \cdot \left[-\frac{1}{5}(x - 2)\right]$   
 $5y + 5 = -(x - 2)$   
 $5y + 5 = -x + 2$   
 $x + 5y = -3$

**80.**  $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{1 - (-5)}{4 - 7} = \frac{6}{-3} = -2$   
 $y - y_1 = m(x - x_1)$   
 $y + 5 = -2(x - 7)$   
 $y + 5 = -2x + 14$   
 $2x + y = 9$

**81.**  $m = 0 \Rightarrow$  horizontal  
 $y = 17$

**82.**  $m$  is undefined  $\Rightarrow$  vertical  
 $x = -5$

**83.**  $3x - 4y = 7$   
 $-4y = -3x + 7$   
 $y = \frac{3}{4}x - \frac{7}{4}$   
 $m = \frac{3}{4}$   
Use  $m = \frac{3}{4}$ .  
 $y - y_1 = m(x - x_1)$   
 $y - 0 = \frac{3}{4}(x - 2)$   
 $y = \frac{3}{4}x - \frac{3}{2}$

**84.**  $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{-10 - 4}{4 - 2} = -7$   
 $y - y_1 = m(x - x_1)$   
 $y + 2 = -7(x - 7)$   
 $y + 2 = -7x + 49$   
 $y = -7x + 47$

## CHAPTER 2 REVIEW

**85.**  $x + 3y = 4$   
 $3y = -x + 4$   
 $y = -\frac{1}{3}x + \frac{4}{3}$   
 $m = -\frac{1}{3}$   
 Use  $m = 3$ .  
 $y - y_1 = m(x - x_1)$   
 $y - 5 = 3(x - 0)$   
 $y - 5 = 3x$   
 $y = 3x + 5$

**87.** Let  $x$  = the number of rolls hung and let  $y$  = the total charge. Then two points on the line are given: (11, 177) and (20, 294)  
 $m = \frac{294 - 177}{20 - 11} = \frac{117}{9} = 13$   
 $y - y_1 = m(x - x_1)$   
 $y - 177 = 13(x - 11)$   
 $y - 177 = 13x - 143$   
 $y = 13x + 34$


Let  $x = 27$ :

$$y = 13(27) + 34 = 385. \text{ The charge is \$385.}$$

**89.**  $y = 4x - 8x^2$        $y = 4x - 8x^2$   
 $0 = 4x(1 - 2x)$        $y = 4(0) - 8(0)^2$   
 $x = 0, x = \frac{1}{2}$        $y = 0$   
 x-int:  $(0, 0), (\frac{1}{2}, 0)$       y-int:  $(0, 0)$

**86.**  $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{-10 - 4}{4 - 2} = -7$   
 Use  $m = \frac{1}{7}$ .  
 $y - y_1 = m(x - x_1)$   
 $y + 2 = \frac{1}{7}(x - 7)$   
 $y + 2 = \frac{1}{7}x - 1$   
 $y = \frac{1}{7}x - 3$

**88.**  $14x + 18y = 5040$   
 Let  $x = 180$ :  
 $14(180) + 18y = 5040$   
 $2520 + 18y = 5040$   
 $18y = 2520$   
 $y = 140$



140 hours of tutoring Spanish

**90.**  $y = x^2 - 10x - 24$        $y = x^2 - 10x - 24$   
 $0 = (x - 12)(x + 2)$        $y = 0^2 - 10(0) - 24$   
 $x = 12, x = -2$        $y = -24$   
 x-int:  $(12, 0), (-2, 0)$       y-int:  $(0, -24)$

**91.**  $y^2 = 8x$

$x$ -axis	$y$ -axis	$y$ -axis
$(-y)^2 = 8x$	$y^2 = 8(-x)$	$(-y)^2 = 8(-x)$
$y^2 = 8x$	$y^2 = -8x$	$y^2 = -8x$
equivalent: <input type="checkbox"/> symmetry	not equivalent: no symmetry	not equivalent: no symmetry

**92.**  $y = 3x^4 + 6$

$x$ -axis	$y$ -axis	$y$ -axis
$-y = 3x^4 + 6$	$y = 3(-x)^4 + 6$	$-y = 3(-x)^4 + 6$
not equivalent: no symmetry	$y = 3x^4 + 6$	$-y = 3x^4 + 6$
	equivalent: <input type="checkbox"/> symmetry	not equivalent: no symmetry

## CHAPTER 2 REVIEW

**93.**

$x\text{-axis}$	$y = -2 x $
$-y = -2 x $	$y = 2 x $

not equivalent: no symmetry

$y = -2 x $	$y\text{-axis}$
-------------	-----------------

$y = -2 -x $	$y = -2 -1  x $
$y = -2 x $	$y = 2 -x $

equivalent: symmetry

$-y = -2 -x $	$\text{origin}$
---------------	-----------------

$y = 2 -x $	$y = 2 -1  x $
$y = 2 x $	$y = 2 x $

not equivalent: no symmetry

**94.**

$x\text{-axis}$	$y =  x + 2 $
$-y =  x + 2 $	$y =  -x + 2 $

not equivalent: no symmetry

$y =  x + 2 $	$y\text{-axis}$
---------------	-----------------


$y =  -x + 2 $	$\text{origin}$
----------------	-----------------

$-y =  -x + 2 $	$-y =  x + 2 $
-----------------	----------------

not equivalent: no symmetry


**95.**  $y = x^2 + 2$

$x\text{-int: none}, y\text{-int: } (0, 2)$   
symmetry:  $y\text{-axis}$




**96.**  $y = -x^2 + 9$

$x\text{-int: } (\pm 3, 0), y\text{-int: } (0, 9)$   
symmetry:  $y\text{-axis}$




**97.**  $y = x^3 - 2$

$x\text{-int: } (\sqrt[3]{2}, 0),$   
 $y\text{-int: } (0, -2)$   
symmetry: none




**98.**  $y = \sqrt{x} + 2$

$x\text{-int: none}, y\text{-int: } (0, 2)$   
symmetry: none




**99.**  $y = -\sqrt{x - 4}$

$x\text{-int: } (4, 0), y\text{-int: none}$   
symmetry: none




**100.**  $y = \frac{1}{2}|x|$

$x\text{-int: } (0, 0), y\text{-int: } (0, 0)$   
symmetry:  $y\text{-axis}$




## CHAPTER 2 REVIEW


**101.**  $y = |x + 1| + 2$   
 x-int: none, y-int: (0, 3)  
 symmetry: none




**102.**  $y = \sqrt[3]{x} - 1$   
 x-int: (1, 0), y-int: (0, -1)  
 symmetry: none




**103.**  $y = |x - 4| + 2$



**104.**  $y = -\sqrt{x+2} + 3$




**105.**  $y = x + 2|x|$



**106.**  $y^2 = x - 3$

Graph  $y = \pm \sqrt{x - 3}$ .



**107.**  $x^2 + y^2 = 64$   
 $(x - 0)^2 + (y - 0)^2 = 8^2$   
 C: (0, 0);  $r = 8$

**108.**  $x^2 + (y - 6)^2 = 100$   
 $(x - 0)^2 + (y - 6)^2 = 10^2$   
 C: (0, 6);  $r = 10$

**109.**  $(x + 7)^2 + y^2 = \frac{1}{4}$   
 $(x - (-7))^2 + (y - 0)^2 = (\frac{1}{2})^2$   
 C: (-7, 0);  $r = \frac{1}{2}$

**110.**  $(x - 5)^2 + (y + 1)^2 = 9$   
 $(x - 5)^2 + (y - (-1))^2 = 3^2$   
 C: (5, -1);  $r = 3$

**111.**  $(x - 0)^2 + (y - 0)^2 = 7^2$   
 $x^2 + y^2 = 49$

**112.**  $(x - 3)^2 + (y - 0)^2 = (\frac{1}{5})^2$   
 $(x - 3)^2 + y^2 = \frac{1}{25}$

**113.**  $(x - (-2))^2 + (y - 12)^2 = 5^2$   
 $(x + 2)^2 + (y - 12)^2 = 25$

**114.**  $(x - \frac{2}{7})^2 + (y - 5)^2 = 9^2$   
 $(x - \frac{2}{7})^2 + (y - 5)^2 = 81$

## CHAPTER 2 REVIEW

**115.**  $C(-3, 4); r = 12$

$$(x - h)^2 + (y - k)^2 = r^2$$

$$(x + 3)^2 + (y - 4)^2 = 144$$

or  $x^2 + y^2 + 6x - 8y - 119 = 0$

**117.**  $x^2 + y^2 + 6x - 4y + 4 = 0$

$$x^2 + 6x + y^2 - 4y = -4$$


$$x^2 + 6x + 9 + y^2 - 4y + 4 = -4 + 9 + 4$$

$$(x + 3)^2 + (y - 2)^2 = 9$$

**119.**  $x^2 + y^2 - 16 = 0$

$$(x - 0)^2 + (y - 0)^2 = 16$$

$C(0, 0), r = 4$




**121.**  $x^2 + y^2 - 2y = 15$

$$x^2 + y^2 - 2y + 1 = 15 + 1$$

$$x^2 + (y - 1)^2 = 16$$

$C(0, 1), r = 4$



**116.** Center:  $x = \frac{-6 + 5}{2} = -\frac{1}{2}$

$$y = \frac{-3 + 8}{2} = \frac{5}{2}$$

$r = \text{distance from center to endpoint}$

$$= \sqrt{\left(-\frac{1}{2} - 5\right)^2 + \left(\frac{5}{2} - 8\right)^2} = \sqrt{\frac{121}{2}}$$

$$\left(x + \frac{1}{2}\right)^2 + \left(y - \frac{5}{2}\right)^2 = \frac{121}{2}, \text{ or}$$

$$x^2 + y^2 + x - 5y - 54 = 0$$

**118.**  $2x^2 + 2y^2 - 8x - 16y - 10 = 0$

$$x^2 + y^2 - 4x - 8y - 5 = 0$$

$$x^2 - 4x + y^2 - 8y = 5$$

$$x^2 - 4x + 4 + y^2 - 8y + 16 = 5 + 4 + 16$$

$$(x - 2)^2 + (y - 4)^2 = 25$$


**120.**  $x^2 + y^2 - 4x = 5$

$$x^2 - 4x + y^2 = 5$$

$$x^2 - 4x + 4 + y^2 = 5 + 4$$

$$(x - 2)^2 + y^2 = 9$$

$C(2, 0), r = 3$




**122.**  $x^2 + y^2 - 4x + 2y = 4$

$$x^2 - 4x + 4 + y^2 + 2y + 1 = 4 + 4 + 1$$

$$(x - 2)^2 + (y + 1)^2 = 9$$

$C(2, -1), r = 3$




## CHAPTER 2 REVIEW

- 123.** Graph  $y = x^2 - 11$ .

Find the  $x$ -intercepts.


$$x = -3.32, x = 3.32$$



- 125.** Graph  $y = |x^2 - 2| - 1$ .

Find the  $x$ -intercepts.

$$x = -1.73, x = -1, x = 1, x = 1.73$$



**127.**  $\frac{x+3}{10} = \frac{x-1}{x}$

$$x(x+3) = 10(x-1)$$

$$x^2 + 3x = 10x - 10$$

$$x^2 - 7x + 10 = 0$$

$$(x-5)(x-2) = 0$$

$$x = 5 \text{ or } x = 2$$

- 129.** Let  $x$  = the dosage needed.

$$\frac{250}{110} = \frac{x}{176}$$

$$250 \cdot 176 = 110 \cdot x$$

$$44000 = 110x$$


$$400 = x$$

The dosage is 400 mg.

- 124.** Graph  $y = x^3 - x$ .

Find the  $x$ -intercepts.


$$x = -1, x = 0, x = 1$$



- 126.** Graph  $y = x^2 - 3x - 5$ .

Find the  $x$ -intercepts.

$$x = -1.19, x = 4.19$$



**128.**  $\frac{x-1}{2} = \frac{12}{x+1}$

$$(x+1)(x-1) = 2(12)$$

$$x^2 - 1 = 24$$

$$x^2 = 25$$

$$x = \pm 5$$

**130.**  $f = ks$        $f = \frac{3}{5}s$   
 $3 = k(5)$        $f = \frac{3}{5}(3)$   
 $\frac{3}{5} = k$        $f = \frac{9}{5}$  pounds

## CHAPTER 2 REVIEW

**131.**  $E = kv^2$   
 30 mph      50 mph  
 $E = k(30)^2$      $E = k(50)^2$   
 $E = 900k$      $E = 2500k$   
 Factor of increase =  $\frac{2500k}{900k} = \frac{25}{9}$

**132.**  $V = \frac{kT}{P}$        $V = \frac{\frac{100}{3}T}{P}$   
 $400 = \frac{k(300)}{25}$      $V = \frac{\frac{100}{3}(200)}{20}$   
 $10000 = 300k$      $V = \frac{1000}{3}$   
 $\frac{100}{3} = k$        $V = 333\frac{1}{3} \text{ cm}^3$

**133.**  $A = klw$   
 $A = 1lw \Rightarrow k = 1$

**134.**  $R = \frac{kL}{D^2}$        $R = \frac{0.0005L}{D^2}$   
 $200 = \frac{k(1000)}{(.05)^2}$      $V = \frac{0.0005(1500)}{(0.08)^2}$   
 $200 = \frac{1000k}{.0025}$      $V \approx 117 \text{ ohms}$   
 $0.0005 = k$

### Chapter 2 Test (page 291)

**1.**  $f(x) = \frac{3}{2x-5}$   
 domain =  $\left(-\infty, \frac{5}{2}\right) \cup \left(\frac{5}{2}, \infty\right)$

**2.**  $f(x) = \sqrt{x+3}$ : domain =  $[-3, \infty)$

**3.**  $f(-1) = \frac{-1}{-1-1} = \frac{-1}{-2} = \frac{1}{2}$   
 $f(2) = \frac{2}{2-1} = \frac{2}{1} = 2$

**4.**  $f(-1) = \sqrt{-1+7} = \sqrt{6}$   
 $f(2) = \sqrt{2+7} = \sqrt{9} = 3$


**5.** 
$$\begin{aligned} \frac{f(x+h)-f(x)}{h} &= \frac{[(x+h)^2 - (x+h)+5] - [x^2 - x+5]}{h} \\ &= \frac{[x^2 + 2xh + h^2 - x - h + 5] - [x^2 - x + 5]}{h} \\ &= \frac{x^2 + 2xh + h^2 - x - h + 5 - x^2 + x - 5}{h} \\ &= \frac{2xh + h^2 - h}{h} = \frac{h(2x + h - 1)}{h} = 2x + h - 1 \end{aligned}$$

**6.**  $(-3, \pi) \Rightarrow \text{QII}$


**7.**  $(0, -8) \Rightarrow \text{negative } y\text{-axis}$

## CHAPTER 2 TEST


8.  $x + 3y = 6$      $x + 3y = 6$   
 $x + 3(0) = 6$      $0 + 3y = 6$   
 $x = 6$                $y = 2$   
 $(6, 0)$                $(0, 2)$




10.  $2(x + y) = 3x + 5$      $\begin{array}{|c|c|} \hline x & y \\ \hline 0 & \frac{5}{2} \\ \hline 1 & 3 \\ \hline \end{array}$   
 $2x + 2y = 3x + 5$   
 $2y = x + 5$   
 $y = \frac{1}{2}x + \frac{5}{2}$




12.  $\frac{1}{2}(x - 2y) = y - 1$      $\begin{array}{|c|c|} \hline x & y \\ \hline 0 & \frac{1}{2} \\ \hline 2 & 1 \\ \hline \end{array}$   
 $\frac{1}{2}x - y = y - 1$   
 $x - 2y = 2y - 2$   
 $-4y = -x - 2$   
 $y = \frac{1}{4}x + \frac{1}{2}$




9.  $2x - 5y = 10$      $2x - 5y = 10$   
 $2x - 5(0) = 10$      $2(0) - 5y = 10$   
 $x = 5$                $y = -2$   
 $(5, 0)$                $(0, -2)$



11.  $3x - 5y = 3(x - 5)$      $\begin{array}{|c|c|} \hline x & y \\ \hline 0 & 3 \\ \hline -2 & 3 \\ \hline \end{array}$   
 $3x - 5y = 3x - 15$   
 $-5y = -15$   
 $y = 3$



13.  $\frac{x + y - 5}{7} = 3x$      $\begin{array}{|c|c|} \hline x & y \\ \hline 0 & 5 \\ \hline -\frac{1}{4} & 0 \\ \hline \end{array}$   
 $x + y - 5 = 21x$   
 $y = 20x + 5$



## CHAPTER 2 TEST

- 14.**  $d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$
- $$\begin{aligned} &= \sqrt{(1 - (-3))^2 + (-1 - 4)^2} \\ &= \sqrt{(4)^2 + (-5)^2} \\ &= \sqrt{16 + 25} = \sqrt{41} \end{aligned}$$
- 15.**  $d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$
- $$\begin{aligned} &= \sqrt{(0 - (-\pi))^2 + (\pi - 0)^2} \\ &= \sqrt{\pi^2 + \pi^2} \\ &= \sqrt{2\pi^2} = \pi\sqrt{2} \approx 4.44 \end{aligned}$$
- 16.**  $M\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}\right) = M\left(\frac{3 + (-3)}{2}, \frac{-7 + 7}{2}\right) = M\left(\frac{0}{2}, \frac{0}{2}\right) = M(0, 0)$
- 17.**  $M\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}\right) = M\left(\frac{0 + \sqrt{8}}{2}, \frac{\sqrt{2} + \sqrt{18}}{2}\right) = M\left(\frac{2\sqrt{2}}{2}, \frac{4\sqrt{2}}{2}\right) = M(\sqrt{2}, 2\sqrt{2})$
- 18.**  $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{1 - (-9)}{-5 - 3} = \frac{10}{-8} = -\frac{5}{4}$
- 19.**  $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{0 - 3}{-\sqrt{12} - \sqrt{3}} = \frac{-3}{-3\sqrt{3}} = \frac{1}{\sqrt{3}} = \frac{\sqrt{3}}{3}$
- 20.**  $y = 3x - 2 \quad y = 2x - 3$   
 $m = 3 \quad m = 2$   
neither
- 21.**  $2x - 3y = 5 \quad 3x + 2y = 7$   
 $-3y = -2x + 5 \quad 2y = -3x + 7$   
 $y = \frac{2}{3}x - \frac{5}{3} \quad y = -\frac{3}{2}x + \frac{7}{2}$   
 $m = \frac{2}{3} \quad m = -\frac{3}{2}$   
perpendicular
- 22.**  $y - y_1 = m(x - x_1)$   
 $y + 5 = 2(x - 3)$   
 $y + 5 = 2x - 6$   
 $y = 2x - 11$
- 23.**  $y = mx + b$   
 $y = 3x + \frac{1}{2}$
- 24.**  $2x - y = 3$   
 $-y = -2x + 3$   
 $y = 2x - 3$   
 $m = 2 \quad y = 2x + 5$
- 25.**  $2x - y = 3$   
 $-y = -2x + 3$   
 $y = 2x - 3$   
 $m = 2 \quad y = -\frac{1}{2}x + 5$
- 26.**  $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{\frac{1}{2} - \left(-\frac{3}{2}\right)}{3 - 2} = \frac{\frac{4}{2}}{1} = 2$   
 $y - y_1 = m(x - x_1)$   
 $y - \frac{1}{2} = 2(x - 3)$   
 $y - \frac{1}{2} = 2x - 6$   
 $y = 2x - \frac{11}{2}$
- 27.** If the line is parallel to the  $y$ -axis, then it is a vertical line:  $x = 3$

## CHAPTER 2 TEST

**28.**  $y = x^3 - 16x$   
 $0 = x(x^2 - 16)$   
 $0 = x(x + 4)(x - 4)$   
 $x = 0, x = -4, x = 4$   
 x-int:  $(0, 0), (-4, 0), (4, 0)$

$$y = x^3 - 16x$$

$$y = 0^3 - 16(0)$$

$$y = 0$$

$$y\text{-int: } (0, 0)$$

**29.**  $y = |x - 4|$   
 $0 = |x - 4|$   
 $0 = x - 4$   
 $4 = x$   
 x-int:  $(4, 0)$

$$y = |x - 4|$$

$$y = |0 - 4|$$

$$y = |-4|$$

$$y = 4$$

**30.**  $y^2 = x - 1$   
 x-axis  
 $(-y)^2 = x - 1$   
 $y^2 = x - 1$   
 equivalent:  symmetry

y-axis  
 $y^2 = -x - 1$   
 not equivalent: no symmetry


origin  
 $(-y)^2 = -x - 1$   
 $y^2 = -x - 1$   
 not equivalent: no symmetry

**31.**  $y = x^4 + 1$   
 x-axis  
 $-y = x^4 + 1$   
 not equivalent: no symmetry


y-axis  
 $y = (-x)^4 + 1$   
 $y = x^4 + 1$   
 equivalent:  symmetry

origin  
 $-y = (-x)^4 + 1$   
 $-y = x^4 + 1$   
 not equivalent: no symmetry


**32.**  $y = x^2 - 9$   
 x-int:  $(3, 0), (-3, 0)$   
 y-int:  $(0, -9)$   
 symmetry: y-axis




**33.**  $x = |y|$   
 x-int:  $(0, 0)$   
 y-int:  $(0, 0)$   
 symmetry: x-axis



**34.**  $y = 2\sqrt{x}$   
 x-int:  $(0, 0)$   
 y-int:  $(0, 0)$   
 symmetry: none



**35.**  $x = y^3$   
 x-int:  $(0, 0)$   
 y-int:  $(0, 0)$   
 symmetry: origin



## CHAPTER 2 TEST


36.  $C(5, 7); r = 8$

$$(x - h)^2 + (y - k)^2 = r^2$$

$$(x - 5)^2 + (y - 7)^2 = 64$$

38.  $x^2 + y^2 = 9$

$$C(0, 0), r = 3$$



40.  $y = kz^2$

42.  $P = kQ \quad P = \frac{7}{2}Q$

$$7 = k(2)$$


$$\frac{7}{2} = k \quad P = \frac{7}{2}(5)$$

$$P = \frac{35}{2}$$

44. Graph  $y = x^2 - 7$ .

Find any positive  $x$ -intercept.

$$x = 2.65$$



37.  $r = \sqrt{(2 - 6)^2 + (4 - 8)^2}$

$$= \sqrt{32}$$

$$(x - h)^2 + (y - k)^2 = r^2$$

$$(x - 2)^2 + (y - 4)^2 = 32$$


39.  $x^2 - 4x + y^2 + 3 = 0$

$$x^2 - 4x + y^2 = -3$$

$$x^2 - 4x + 4 + y^2 = -3 + 4$$

$$(x - 2)^2 + y^2 = 1$$

$$C(2, 0), r = 1$$



41.  $w = krs^2$

43.  $y = \frac{kx}{z^2} \quad y = \frac{\frac{64}{3}x}{z^2}$

$$16 = \frac{k(3)}{2^2} \quad 2 = \frac{\frac{64}{3}x}{3^2}$$

$$16 = \frac{3k}{4} \quad 18 = \frac{64}{3}x$$

$$\frac{64}{3} = k \quad \frac{3}{64} \cdot 18 = \frac{3}{64} \cdot \frac{64}{3}x$$

$$\frac{27}{32} = x$$

45. Graph  $y = x^2 - 5x - 5$ .

Find any positive  $x$ -intercept.

$$x = 5.85$$

